


British Lichen Society
Bulletin No. 72 (Supplement) Summer 1993

CHECKLIST OF LICHENS OF GREAT BRITAIN AND IRELAND



O.W. Purvis, B.J. Coppins and P.W. James

CHECKLIST OF LICHENS OF GREAT BRITAIN AND IRELAND

Completed March 1993

O.W. Purvis*, B.J. Coppins⁺ & P.W. James*

* Department of Botany, The Natural History Museum, Cromwell Rd,
London SW7 5BD

+ The Herbarium, Royal Botanic Garden, Inverleith Row, Edinburgh
EH3 5LR

*For further details of the British Lichen Society contact Dr O.W. Purvis, The
Secretary, The Natural History Museum, Cromwell Rd, London SW7 5BD.
Tel. 071-938 8852, Fax 071-938 9260.*

Introduction

This checklist of the lichens of Great Britain and Ireland includes all accepted names of lichen-forming fungi treated in the *Lichen Flora of Great Britain and Ireland* (Purvis *et al.* 1992). Lichenicolous fungi (suffixed "[LF]") included are those present in genera mainly comprising lichenized species (e.g. *Arthonia* and *Opegrapha* and suffixed "[±L]") and a few that until recently were included in such genera (e.g. *Geltingia* and *Phaeopyxis* species, previously in *Lecidea*). A new checklist of lichenicolous taxa is in preparation. Other non-lichenized taxa (suffixed "[F]"), also included in the *Flora*, are those traditionally treated by lichenologists and which are usually overlooked by mycologists. Earlier synonyms used in the previous checklists of Hawksworth *et al.* (1980) and Cannon *et al.* (1985) are also included. Some later synonyms, especially for recent generic segregates used by some contemporary lichenologists which it has not been prudent to adopt at this stage, are also noted. 22 recently discovered species are also included and listed separately (Appendix 3); these will be discussed in a future paper in the *Lichenologist*. 50 lichen taxa which have not been recorded during this century and which may be regarded as extinct are indicated in the main list by an asterisk (*) and also listed in Appendix 1; species which are considered to have been erroneously reported from Great Britain and Ireland and those which are of dubious taxonomic validity and have been not dealt with in the *Flora* are omitted from the main list and are presented in Appendix 2. New combinations are given in Appendix 4: A "?" denotes in the case of species an uncertain taxon in Great Britain, or, in the case of synonyms a probable synonym.

The present checklist is arranged alphabetically by genus and species. Accepted genera are printed in bold and those regarded as synonyms in bold italics. Accepted species names (with dates of valid publication) are indented in roman, with synonyms further indented in italics. Synonyms which have been transferred to other genera are listed separately at the base of each genus listing. References to literature have been kept to a minimum as these are given in the *Flora* which should be consulted for further information. The total number of taxa accepted in the checklist is 267 genera, 1660 species, 9 subspecies, 40 varieties and 12 forms. Of these 1566 species are consistently or facultatively lichenized; 42 are non-lichenized (but not lichenicolous); and 52 are lichenicolous fungi (excluding lichenicolous lichens). An equivalent 223 lichen genera were included in the 1980 list and 1471 lichen-forming fungi were listed; the difference between these figures and the current totals reflects the discovery of a substantial number of additional species in the British Isles as well as a proliferation in genera, particularly as a consequence of the splitting of *Lecidea* and other crustose genera.

Checklist

Absconditella Vězda (1965)

- celata Döbbeler & Poelt (1977)
- delutula (Nyl.) Coppins & Kilias (1981)
- ? *A. modesta* (Stizenb.) Vězda (1965)
- pauxilla Vězda & Vivant (1975)
- sphagnorum Vězda & Poelt (1965)
- trivialis (Willey ex Tuck.) Vězda (1965)

Acarospora Massal. (1852)

- badiofusca (Nyl.) Th. Fr. (1861)
- chlorophhana (Wahlenb.) Massal. (1852)
- durietzii Magnusson (1924)
- fuscata (Schrader) Th. Fr. (1871)
- glaucocarpa (Ach.) Körber (1859)
- ? *A. cervina* Massal. (1852)
- A. glaucocarpa* var. *depauperata* (Körber) A.L. Sm. (1918)
- heppii (Naeg. ex Hepp) Naeg. ex Körber (1859)
- impressula Th. Fr. (1871)
- A. atrata* Hue (1909)
- macrospora (Hepp) Massal. ex Bagl. (1857)
- A. macrospora* var. *incusa* (Körber) Magnusson (1924)
- macrospora subsp. *muronorum* (Massal.) Clauz. & Roux (1981)
- A. murorum* Massal. (1853)
- nitrophila Magnusson (1924)
- A. muddii* Magnusson (1926)
- A. normanii* Magnusson (1924)
- A. praeruptorum* Magnusson (1924)
- rhizobola (Nyl.) Alstrup (1986)
- Lecidea rhizobola* Nyl. (1865)
- rufescens (Ach.) Krempelh. (1861)
- sinopica (Wahlenb.) Körber (1859)
- smaragdula (Wahlenb.) Massal. (1852)
- A. amphibola* Wedd. (1875)
- A. benedarensis* Knowles (1913)
- A. fusca* B. de Lesd. (1914)
- A. scabrida* auct. brit. p.p., non Hedl. ex Magnusson (1924)
- A. scyphulifera* Vainio (1909)
- A. smaragdula* var. *lesdainii* (Harm. ex A.L. Sm.) Magnusson (1929)
- A. smaragdula* var. *murina* (Sandst.) Magnusson (1929)
- subrufula (Nyl.) H. Olivier (1900)
- umbilicata Bagl. (1857) f. *congregans* Magnusson (1929)
- veronensis Massal. (1852)
- A. magnussonii* G. Samp. (1924)
- verruciformis Magnusson (1924)

Acrocordia Massal. (1854)

- cavata (Ach.) R. Harris (1974)
conoidea (Fr.) Körber (1855)
gemma (Ach.) Massal. (1854)
macrospora Massal. (1855)
salweyi (Leighton ex Nyl.) A.L. Sm. (1911)

Adelolecia Hertel & Hafellner (1984)

- pilati (Hepp) Hertel & Hafellner (1984)
Lecidea pilati (Hepp) Körber (1861)

Agonimia Zahlbr. (1909)

- allobata (Stizenb.) P. James (1992)
Polyblastia allobata (Stizenb.) Zsch. (1918)
octospora Coppins & P. James (1978)
tristicula (Nyl.) Zahlbr. (1909)

Agyrium Fr. (1822) [F]

- rufum (Pers.) Fr. (1822) [F]
Lecidea grumosa Leighton (1876)

Ahlesia Fuckel (1870) - see *Thelocarpon*.

Alectoria Ach. (1810)

- nigricans (Ach.) Nyl. (1861)
ochroleuca (Hoffm.) Massal. (1855)
sarmentosa (Ach.) Ach. (1810)
sarmentosa subsp. vexillifera (Nyl.) D. Hawksw. (1970)

Allantoparmelia (Vainio) Essl. (1978)

- alpicola (Th. Fr.) Essl. (1978)

Amygdalaria Norman (1853)

- consentiens (Nyl.) Hertel, Brodo & Inoue (1984)
Hulia consentiens (Nyl.) Hertel (1980)
pelobotryon (Wahlenb.) Norman (1953)

Anaptychia Körber (1848)

- ciliaris Körber ex Massal. (1853)
ciliaris subsp. mamillata (Taylor) D. Hawksw. & P. James (1980)
runcinata (With.) Laundon (1984)
A. fusca (Huds.) Vainio (1899)

Anisomeridium (Müll. Arg.) M. Choisy (1928)
 biforme (Borrer) R. Harris (1978)
 nyssaenum (Ellis & Everh.) R. Harris (1985)
A. juistense (Erichsen) R. Harris (1980)
Sarcinulella banksiae B. Sutton & Alcorn (1983)

Aphanopsis Nyl. ex Sydow (1887)
coenosa (Ach.) Coppins & P. James (1984)
Lecidea humigena Taylor (1847)

Arctomia Th. Fr. (1860)
delicatula Th. Fr. (1860)

Arctoparmelia Hale (1986) - see *Parmelia*.

Arthonia Ach. (1806) [±L]
abelonea P.M. Jørg. (1970) [LF]
amylospora Almq. (1880) [LF]
anglica Coppins (1989)
anombrophila Coppins & P. James (1989)
A. cinereopruinosa auct. brit., non Schaerer (1850)
arthonioides (Ach.) A.L. Sm. (1911)
A. aspersa Leighton (1856)
astroidestra Nyl. (1874)
A. stellaris auct. p.p. non Krempelh. (1861)
A. zwackhii var. *macrospora* P. James (1978)
atlantica P. James (1970)
A. atlantica var. *positiva* P. James (1978)
cinnabarina (DC.) Wallr. (1831)
A. tumidula (Ach.) Ach. (1806)
clemens (Tul.) Th. Fr. (1867) [LF]
cohabitans Coppins (1989) [LF]
didyma Körber (1853)
A. aspersella Leighton (1872)
elegans (Ach.) Almq. (1880)
endlicheri (Garov.) Oxner (1956)
epiphyscia Nyl. (1875) [LF]
excipienda (Nyl.) Leighton (1871) [LF]
A. dispersa auct. brit. p.p., non (Schrader) Nyl. (1861)
A. hibernica Nyl. (1876)
fuscopurpurea (Tul.) R. Sant. (1960) [LF]
* *galactites* (DC.) Dufour (1818)
gelidae R. Sant. (1986) [LF]
graphidicola Coppins (1869) [LF]
ilicina Taylor (1836)
Arthothelium ilicinum (Taylor) P. James (1965)
ilicinella Nyl. (1867)
impolitata (Hoffm.) Borrer (1831)

- intexta Almq. (1880) [LF]
 invadens Coppins (1989) [LF]
 lapidicola (Taylor) Branth. & Rostrup (1869)
Catillaria ooliticola W. Watson (1935)
 leucopellaea (Ach.) Almq. (1880)
 lignaria Hellbom (1884)
 lignariella Coppins (1991)
 ? molendoi (Heufl. ex Frauendorf.) R. Sant. (1986) [LF]
 muscigena Th. Fr. (1865)
 A. exilis auct. brit., non (Flörke) Anzi (1860)
 A. leucodontis (Poelt & Döbbeler) Coppins (1989)
 Bryostigma leucodontis Poelt & Döbbeler (1979)
 * myriocarrella Nyl. (1863)
 peltigerae Th. Fr. (1866) [LF]
 phaeobaea (Norman) Norman (1869)
 punctella Nyl. (1859) [LF]
 punctiformis Ach. (1808) [LF]
 radiata (Pers.) Ach. (1808)
 spadicea Leighton (1854)
 stellaris Krempelh. (1861)
 * subfuscicola (Lindsay) Triebel (1991) [LF]
 thelotrematis Coppins (1989) [LF]
 varians (Davies) Nyl. (1861) [LF]
 A. glaucomaria (Nyl.) Nyl. (1861)
 vinosa Leighton (1856)
 zwackhii Sandst. (1903)

A. granitophila Th. Fr. (1865) = *Melaspilea*.

- Arthopyrenia** Massal. (1852). [\pm L]
 antecellans (Nyl.) Arnold (1870) [F]
 carneobrunnea Coppins (1988)
 cerasi (Schrader) Massal. (1852) [F]
 A. crombei A.L. Sm. (1911)
 cinereopruinosa (Schaerer) Massal. (1855) [F]
 fraxini Massal. (1852) [F]
 lapponina Anzi (1864) [F]
 nitescens (Salwey) Mudd (1861)
 A. submicens (Nyl.) Arnold (1873)
 punctiformis Massal. (1852) [F]
 A. cembra (Anzi) Grumman ex D. Hawksw. (1980)
 A. laburni Arnold (1861)
 ? *A. padi* Rabenh. (1858)
 ranunculospora Coppins & P. James (1988)
 rhyponta (Ach.) Massal. (1852) [F]
 salicis Massal. (1852)
 subcerasi (Vainio) Zahlbr. (1921) [F]
 viridescens Coppins (1988)

A. areniseda A.L. Sm. (1911), *A. argilospora* (Nyl.) A.L. Sm. (1918), *A. bryospila* (Nyl.) Arnold (1870), *A. caesia* (Nyl.) Zahlbr. (1921), *A. elegans* R. Sant, nom. inval. (1985), *A. halodites* (Nyl.) Arnold (1891), *A. monensis* (Wheldon) Zahlbr. (1922), *A. orustensis* Erichsen (1930), *A. pelvetiae* (Suterl.) D. Hawksw. (1980), *A. saxicola* Massal. (1865), *A. strontianensis* Swinscow (1967), *A. subareniseda* G. Salisb. (1953) and *A. sublitoralis* (Leighton) Arnold (1981) are included in *Pyrenocollema*.

Arthothelium Massal. (1852) [\pm L]

- dictyosporum (Coppins & P. James) Coppins (1989)
- A. ilicina* var. *dictyosporum* Coppins & P. James (1979)
- lirellans (Almq.) Coppins (1979) [F]
- norvegicum Coppins & Tønsb. (1984)
- orbilliferum (Almq.) Hasse (1913) [F]
- reagens (Coppins & P. James) Coppins & P. James (1981)
- A. ilicina* var. *reagens* Coppins & P. James (1979)
- ruanum (Massal.) Körber (1861)
- * spectabile Flotow ex Massal. (1852)

A. ilicina (Taylor) P. James (1965) = *Arthonia*.

Arthroraphis Th. Fr. (1860) [\pm L]

- alpina (Schaeerer) R. Sant. (1980)
- citrinella (Ach.) Poelt (1969)
- fuscireagens (Vainio) Poelt (1969) [LF]
- grisea Th. Fr. (1860) [LF]

Aspicilia Massal. (1852)

- caesiocinerea (Nyl. ex Malbr.) Arnold (1886)
- A. gibbosa* auct. brit., p.p. non (Ach.) Körber (1855)
- calcarea (L.) Mudd (1861)
- ? *A. lilliei* B. de Lesd. (1906)
- cinerea (L.) Körber (1855)
- contorta (Hoffm.) Krempelh. (1861)
- epiglypta (Norrlin ex Nyl.) Hue (1912)
- grisea Arnold (1891)
- A. insolata* (Magnusson) Havaas (1944)
- intermutans (Nyl.) Arnold (1887)
- laevata (Ach.) Arnold (1887)
- ? *A. sylvatica* Arnold (1884)
- leprosescens (Sandst.) Havaas (1918)
- melanaspis (Ach.) Poelt & Leuckert (1973)
- recedens (Taylor) Arnold (1896)
- ? simoensis Räsänen (1925)
- subcircinata (Nyl.) Coppins (1980)
- subdepressa auct. brit., non (Nyl.) Arnold (1869)
- * tuberculosa (Ach.) Laundon (1986)

- A. alpina* (Sommerf.) Arnold (1871) = *Bellemerea*
A. complanata (Körber) B. Stein (1879) = *Miriquidica*
A. complanatoides (A.L. Sm.) W. Watson (1935) = *Schaereria*
A. flava (Hepp) Rehm (1863) = *Eiglera*
A. leucophyma (Leighton) Hue (1912) = *Koerberiella*
A. lindsayi W. Watson (1939) = *Lecanora*
A. morioides Blomb. ex Arnold (1881) = *Clauzadeana*
A. poriniformis (Nyl.) Arnold (1870) = *Pertusaria*
A. superiuscula (Nyl.) Hue (1912) = *Miriquidica*

Bacidia de Not. (1846)

- absistens (Nyl.) Arnold (1870)
arceutina (Ach.) Arnold (1869)
B. leightonianana (Larbal. ex Leighton) H. Olivier (1911)
arnoldiana Körber (1860)
assulata (Körber) Vězda (1967)
auerswaldii (Hepp ex Stizenb.) Mig. (1929)
bagliettoana (Massal. & de Not.) Jatta (1900)
B. muscorum (Swartz) Mudd (1861)
beckhausii Körber (1860)
biatorina (Körber) Vainio (1922)
caligans (Nyl.) A. L. Sm. (1911)
carneoglaucha (Nyl.) A. L. Sm. (1911)
chloroticula (Nyl.) A. L. Sm. (1911)
circumspecta (Norrlin & Nyl.) Malme (1895)
delicata (Larbal. ex Leighton) Coppins (1980)
egenula (Nyl.) Arnold (1870)
friesiana (Hepp) Körber (1860)
fuscoviridis (Anzi) Lettau (1912)
hegetschweileri (Hepp) Vainio (1922)
herbarum (Stizenb.) Arnold (1865)
igniari (Nyl.) Oxner (1968)
B. abbrevians (Nyl.) Th. Fr. (1874)
incompta (Borrer ex Hooker) Anzi (1860)
inundata (Fr.) Körber (1855)
laurocerasi (Delise ex Duby) Zahlbr. (1926)
myrtillicola Erichsen (1939)
naegelii (Hepp) Zahlbr. (1909)
phacodes Körber (1860)
* polychroa (Th. Fr.) Körber (1860)
rubella (Hoffm.) Massal. (1852)
sabuletorum (Schreber) Lettau (1912)
B. accedens (Arnold) Lettau (1912)
saxenii Erichsen (1941)
scopulicola (Nyl.) A. L. Sm. (1911)
subcircumspecta Coppins (1992)

- subfuscula* (Nyl.) Th. Fr. (1867)
B. circumpallens (Nyl.) Arnold (1870)
subincompta (Nyl.) Arnold (1870)
* *subturgidula* (Nyl.) Zahlbr. (1926)
trachona (Ach.) Lettau (1912)
vezdae Coppins & P. James (1978)
Fellhanera vezdae (Coppins & P. James) V. Wirth (1987)
viridescens (Massal.) Norman (1868)
viridifarinosa Coppins & P. James (1992)

- B. carneoalbens* (Nyl.) A.L. Sm. (1911) = *Micarea*
B. carneoalbida (Müll. Arg.) Coppins (1988) [= *carneopallida*] = *Biatora*
B. cuprea (Massal.) Lettau (1912) = *Lecania*
B. epixanthoides (Nyl.) Lettau (1912) = *Biatora*
B. fusca (Massal.) Du Rietz (1929) = *Biatora*
B. globulosa (Flörke) Hafellner & V. Wirth (1987) = *Catillaria*
B. obscurata (Sommerf.) Zahlbr. (1905) = *Biatora*
B. plumbina (Anzi) R. Sant. (1960) = *Toninia*
B. subtilis Vězda (1961) = *Fellhanera*
B. tetramera (de Not.) Coppins (1988) = *Biatora*.

- Bactrospora** Massal. (1852)
corticola (Fr.) Almq. (1869)
dryina (Ach.) Massal. (1852)
homalotropa (Nyl.) Egea & Torrente (1989)
Lecanactis homalotropa (Nyl.) Gilenstam (1969)

Literature: Torrente & Egea (1989), Egea & Torrente (1993)

- Baeomyces** Pers. (1794)
placophyllus Ach. (1803)
roseus Pers. (1794)
rufus (Huds.) Rebent. (1804)

Bagliettoa Massal. (1853) - see *Verrucaria*.

- Bellemerea** Hafellner & Roux (1984)
alpina (Sommerf.) Clauz. & Roux. (1984)
Aspicilia alpina (Sommerf.) Arnold (1871)

- Belonia** Körber ex Nyl. (1852)
calcicola W. Watson (1935)
incarnata Th. Fr. & Graewe ex Th. Fr. (1865)
nidarosiensis (Kindt) P.M. Jørg & Vězda (1983)
Clathroporina calcarea W. Watson (1925)
russula Körber ex Nyl. (1857)

Biatora Fr. (1817)

carneoalbida (Müll. Arg.) Coppins (1992)

Bacidia carneoalbida (Müll. Arg.) Coppins (1988) ("carneopallida")

Biatora sphaerooides auct. p.p., non (Dickson) Körber (1853)

* *cuprea* (Sommerf.) Fr. (1831)

Lecidea cuprea Sommerf. (1826)

epixanthoides (Nyl.) Diederich (1989)

Bacidia epixanthoides (Nyl.) Lettau (1912)

epixanthoidiza (Nyl.) Räsänen (1939)

Lecidea efflorescens (Hedl.) Erichsen (1932)

Lecidea epixanthoidiza Nyl. (1875), non auct. brit.

gyrophorica (Tönsb.) Coppins (1993)

Lecidea gyrophorica Tönsb. (1992)

Lecidea epixanthoidiza auct., non Nyl. (1875)

Biatora epixanthoidiza auct., non (Nyl.) Räsänen (1939)

sphaerooides (Dickson) Körber (1853)

Catillaria sphaerooides (Massal.) Schuler (1902)

tetramera (de Not.) Coppins (1992)

Bacidia fusca (Massal.) Du Rietz (1929)

Bacidia obscurata (Sommerf.) Zahlbr. (1905)

Bacidia tetramera (de Not.) Coppins (1988)

Mycobilimbia obscurata (Sommerf.) Rehm (1890)

vernalis (L.) Fr. (1831)

Lecidea vernalis (L.) Ach. (1803)

B. humida Kulh. (1871) = *Moelleropsis*.

Biatorella de Not. (1846)

fossarum (Dufour) Th. Fr. (1874)

hemisphaerica Anzi (1860)

B. fossarum auct. brit., non (Dufour) Th. Fr. (1874)

monasteriensis (Lahm) Lahm (1883)

Biatorina Massal. (1852) - see *Catillaria*.

Blarneya D. Hawksw., Coppins & P. James (1980)

hibernica D. Hawksw., Coppins & P. James (1980)

Bombyliospora de Not. ex Massal. (1852) - see *Megalospora*.

Brigantiae Trevisan (1853)

fuscolutea (Dickson) R. Sant. (1981)

Bryonora Poelt (1983)

curvescens (Mudd) Poelt (1983)

Lecanora curvescens (Mudd) Nyl. (1865)

Bryophagus Nitschke ex Arnold (1862)
gloecapsa Nitschke ex Arnold (1862)

Bryoria Brodo & D. Hawksw. (1977)
bicolor (Ehrh.) Brodo & D. Hawksw. (1977)
capillaris (Ach.) Brodo & D. Hawksw. (1977)
chalybeiformis (L.) Brodo & D. Hawksw. (1977)
furcellata (Fr.) Brodo & D. Hawksw. (1977)
fuscescens (Gyelnik) Brodo & D. Hawksw. (1977)
fuscescens var. *positiva* (Gyelnik) Brodo & D. Hawksw. (1977)
* *implexa* (Hoffm.) Brodo & D. Hawksw. (1977)
B. pseudofuscescens (Gyelnik) Brodo & D. Hawksw. (1977)
lanestris (Ach.) Brodo & D. Hawksw. (1977)
nadvornikiana (Gyelnik) Brodo & D. Hawksw. (1977)
smithii (Du Rietz) Brodo & D. Hawksw. (1977)
subcana (Nyl. ex Stitzenb.) Brodo & D. Hawksw. (1977)

Bryostigma Döbbeler & Poelt (1979) - see *Arthonia*.

Buellia de Not. (1846)
abstracta (Nyl.) H. Olivier (1903)
aethalea (Ach.) Th. Fr. (1874)
B. soraria Th. Fr. (1874)
arborea Coppins & Tønsb. (1992)
arnoldii Servít (1931)
asterella Poelt & M. Sulzer (1974)
badia (Fr.) Massal. (1853)
coniops (Wahlenb. ex Ach.) Th. Fr. (1861)
disciformis (Fr.) Mudd (1861)
erubescens Arnold (1873)
excelsa (Leighton) A.L. Sm. (1911)
B. punctata var. *excelsa* (Leighton) Sheard (1964)
griseovirens (Turner & Borrer ex Sm.) Almb. (1952)
insignis (Naeg. ex Hepp) Th. Fr. (1860)
leptocline (Flotow) Massal. (1854)
leptoclinoides (Nyl.) Steiner (1907)
Hafellia leptoclinoides (Nyl.) Scheid. & Mayrhofer (1986)
meiosperma (Nyl.) Müll. Arg. (1887)
ocellata (Flotow) Körber (1855)
B. verruculosa auct., non (Sm.) Mudd (1861)
pulvrea Coppins & P. James (1978)
pulverulenta (Anzi) Jatta (1900)
punctata (Hoffm.) Massal. (1852)
sanguinolenta Schauer (1965)
saxorum Massal. (1852)
schaereri de Not. (1846)
spuria (Schaerer) Anzi (1860)
stellulata (Taylor) Mudd (1861)

subdisciformis (Leighton) Vainio (1890)
uberior Anzi (1866)
violaceofusca Thor & Mhur (1991)

B. aspersa (Borrer) P. James (1965) = *Rinodina*
B. geophila auct. brit., non (Flörke ex Sommerf.) Lynge (1937) = *Diplotomma*.

Byssoloma Trevisan (1853)

leucoblepharum (Nyl.) Vainio (1926)
marginatum (Arnold) Sérusiaux (1991)
subdiscordans (Nyl.) P. James (1971)

Calicium Pers. (1794)

abietinum Pers. (1797)
adspersum Pers. (1798)
corynellum (Ach.) Ach. (1803)
diploellum Nyl. (1868)
glaucellum Ach. (1803)
lenticulare Ach. (1816)
C. subquercinum Asah. (1932)
parvum Tibell (1975)
* *quercinum* Pers. (1797)
salicinum Pers. (1794)
* *trabinellum* (Ach.) Ach. (1803)
viride Pers. (1794)

Caloplaca Th. Fr. (1860)

alociza (Massal.) Migula (1925)
approximata (Lynge) Magnusson (1946)
aractina (Fr.) Häyrén (1914)
C. fuscoatra (Decuill.) Zahlbr. (1894)
arenaria (Pers.) Müll. Arg. (1862)
C. subpallida Magnusson (1945)
arnoldii (Wedd.) Zahlbr. ex Ginzb. (1915)
atroflava (Turner) Monguillon (1914)
? *C. atroflava* var. *submersa* (Nyl.) Magnusson (1944)
aurantia (Pers.) Steiner (1896)
britannica R. Sant. (1992)
ceracea Laundon (1992)
C. caesiorufa auct. brit., non (Schrader) Flagey (1895)
cerina (Ehrh. ex Hedwig) Th. Fr. (1860)
C. stillicidiorum auct. brit., ? non (Vahl) Lynge (1921)
cerinella (Nyl.) Flagey (1896)
chalybaea (Fr.) Müll. Arg. (1862)
chrysophthalma Degel. (1944)
cinnamomea (Th. Fr.) H. Olivier (1909)
? *C. ammiospila* (Wahlenb. ex Ach.) H. Olivier (1909)
cirrochroa (Ach.) Th. Fr. (1871)

- citrina* (Hoffm.) Th. Fr. (1860)
C. citrina f. *phlogina* (Ach.) D. Hawksw. (1969)
concilians (Nyl.) H. Olivier (1909)
crenularia (With.) Laundon (1984)
C. festiva (Ach.) Zwackh (1864)
* *crenulatella* (Nyl.) H. Olivier (1909)
dalmatica (Massal.) H. Olivier (1909)
C. schaeereri (Arnold) Zahlbr. (1890)
C. velana (Massal.) Du Rietz (1925)
decipiens (Arnold) Blomb. & Forss. (1880)
ferruginea (Huds.) Th. Fr. (1860)
flavescens (Huds.) Laundon (1984)
C. heppiana (Müll. Arg.) Zahlbr. (1890)
flavorubescens (Huds.) Laundon (1976)
flavovirescens (Wulfen) Dalla Torre & Sarnth. (1902)
granulosa (Müll. Arg.) Jatta (1900)
* *haematites* (Chaub. ex St-Amans) Zwackh (1862)
herbidella (Hue) Magnusson (1932)
holocarpa (Hoffm.) Wade (1965)
C. furfuracea sensu Wade (1965), non Magnusson (1944)
* *irrubescens* (Arnold) Zahlbr. (1898)
isidiigera Vězda (1978)
C. cerina var. *cyanopolia* (Nyl.) Zahlbr. (1930)
? *C. chlorina* (Flotow) H. Olivier (1909)
lactea (Massal.) Zahlbr. (1901)
littorea Tavares (1956)
lucifuga Thor (1988)
luteoalba (Turner) Th. Fr. (1860)
marina (Wedd.) Zahlbr. ex Du Rietz (1921)
microthallina (Wedd.) Zahlbr. (1931)
nivalis (Körber) Th. Fr. (1871)
obliterans (Nyl.) Blomb & Forss. (1880)
obscurella (Lahm ex Körber) Th. Fr. (1871)
C. sarcopisioides (Körber) Zahlbr. (1901)
ochracea (Schaerer) Flagey (1886)
* *pollinii* (Massal.) Jatta (1900)
ruderum (Malbr.) Laundon (1976)
saxicola (Hoffm.) Nordin (1972)
C. miniatula (Nyl.) Zahlbr. (1931)
scopularis (Nyl.) Lettau (1912)
teicholyta (Ach.) Steiner (1895)
C. albolutescens (Nyl.) H. Olivier (1909)
C. erythrocarpa auct., non (Pers.) Zwackh (1862)
thallincola (Wedd.) Du Rietz (1921)
ulcerosa Coppins & P. James (1979)
variabilis (Pers.) Müll. Arg. (1862)
verruculifera (Vainio) Zahlbr. (1931)
virescens (Sm.) Coppins (1980)

C. chrysodeta (Vainio ex Räsänen) Dombr. (1970) and *C. xantholyta* (Nyl.) Jatta (1902) are included in *Leproplaca*.

Candelaria Massal. (1852)
concolor (Dickson) B. Stein (1879)

Candelariella Müll. Arg. (1894)
aurella (Hoffm.) Zahlbr. (1928)
aurella f. *smaragdula* Szat. (1931)
C. aurella f. *heidelbergensis* (Nyl.) P. James (1981)
coralliza (Nyl.) Magnusson (1935)
medians (Nyl.) A.L. Sm. (1918)
medians f. *steepholtiensis* O. Gilbert (1981)
reflexa (Nyl.) Lettau (1912)
vitellina (Hoffm.) Müll. Arg. (1894)
vitellina f. *flavovirella* (Nyl.) A. Henderson (1981)
C. flavovirella (Nyl.) Lettau (1912)
xanthostigma (Ach.) Lettau (1912)

Carbonea (Hertel) Hertel (1983) [\pm L]
assimilis (Körber) Hafellner & Hertel (1987)
supersparsa (Nyl.) Hertel (1983) [LF]
Lecidea supersparsa Nyl. (1865)
vitellinaria (Nyl.) Hertel (1983) [LF]
Lecidea vitellinaria Nyl. (1852)
vorticosa (Flörke) Hertel (1983)
Lecidea vorticosa (Flörke) Körber (1855)

Catapyrenium Flotow (1850)
boccanum (Servít) Breuss (1990)
cinereum (Pers.) Körber (1855)
daedaleum (Kremphelh.) B. Stein (1879)
lachneum (Ach.) R. Sant. (1980)
michelli (Massal.) R. Sant. (1980)
pilosellum Breuss (1990)
psoromoides (Borrer) R. Sant. (1980)
rufescens (Ach.) Breuss (1985)
squamulosum (Ach.) Breuss (1985)
Dermatocarpon hepaticum auct., non (Ach.) Th. Fr. (1860)
waltheri (Kremphelh.) Körber (1855)

Catillaria Massal. (1852)
aphana (Nyl.) Coppins (1989)
Lecidea aphana Nyl. (1867)
atomariooides (Müll. Arg.) Kilias (1981)
atropurpurea (Schaerer) Th. Fr. (1874)
Biatorina atropurpurea (Schaerer) Massal. (1852)
Catinaria atropurpurea (Schaerer) Vězda & Poelt. (1981)

- chalybeia* (Borrer) Massal. (1852)
chalybeia var. *chloropoliza* (Nyl.) Kilias (1981)
contristans (Nyl.) Zahlbr. (1926)
globulosa (Flörke) Th. Fr. (1874)
Bacidia globulosa (Flörke) Hafellner & V. Wirth (1987)
laureri Hepp ex Th. Fr. (1874)
Catinaria laureri (Hepp ex Th. Fr.) Degel. (1941)
lenticularis (Ach.) Th. Fr. (1874)
Lecania actea (Nyl.) B. de Lesd. (1910)
* *minuta* (Massal.) Lettau (1912)
modesta (Müll. Arg.) Coppins (1989)
Lecidea modesta Müll. Arg. (1871)
neuschildii (Körber) Th. Fr. (1874)
Biatorina neuschildii Körber (1860)
Catinaria neuschildii (Körber) P. James (1965)
nigroclavata (Nyl.) Schuler (1902)
* *picila* (Massal.) Coppins (1989)
pulvrea (Borrer) Lettau (1912)
Catinaria pulvrea (Borrer) Vězda & Poelt (1981)
scotinodes (Nyl.) Coppins (1989)
C. confusior (Nyl.) Zahlbr. (1926)
Kiliasia scotinodes (Nyl.) Coppins (1988)
Lecidea confusior Nyl. (1874)
Lecidea scotinodes Nyl. (1873)
subviridis (Nyl.) Zahlbr. (1926)

- C. biformigera* (Leighton) Magnusson (1949) = *Tylothallia*
C. bouteillei (Desm.) Zahlbr. (1902) = *Fellhanera*
C. chlorotiza (Nyl.) P. James (1965) = *Lecania*
C. episema (Nyl.) H. Olivier (1905) = *Toninia*
C. intrusa (Th. Fr.) Th. Fr. (1874) = *Micarea*
C. jejuna (Nyl.) P. James (1965) = *Halecania*
C. littorella auct. brit., non (Nyl.) Zahlbr. (1926) = *Lecania*
C. littorella (Nyl.) Zahlbr. (1926), non auct. = *Fellhanera*
C. ooliticola W. Watson (1935) = *Arthonia*
C. rhypodiza (Nyl.) Zahlbr. (1926) = *Halecania*
C. sphaeroides (Massal.) Schuler (1902) = *Biatoria*
C. subnigrata (Nyl.) Herre (1910) = *Micarea*
C. umbraticula (Nyl.) P. James (1965) = *Lecania*.

Catinaria Vainio (1922) - see *Catillaria*.

For *Catinaria grossa* (Pers. ex Nyl.) Vainio (1922) see *Megalaria*.

Catolechia Flotow (1850)
wahlenbergii (Ach.) Körber (1855)

Cavernularia Degel. (1938)
hultenii Degel. (1938)

Cecidonia Triebel & Rambold (1988) [LF]
umbonella (Nyl.) Triebel & Rambold (1988) [LF]
Lecidea umbonella Nyl. (1866)
Lecidea umbonata auct. brit., non (Hepp) Mudd (1861)
xenophana (Körber) Triebel & Rambold (1988) [LF]
Lecidea umbonella var. *alumnula* (Nyl.) Hertel (1970)

Celothelium Massal. (1860)
ischnobelum (Nyl.) M.B. Aguirre (1991)
Leptorraphis ischnobela (Nyl.) Coppins (1980)

Cetraria Ach. (1803)
chlorophylla (Willd.) Vainio (1896)
commixta (Nyl.) Th. Fr. (1871)
delisei (Bory ex Schaeer) Nyl. (1866)
ericetorum Opiz (1852)
hepatizon (Ach.) Vainio (1899)
islandica (L.) Ach. (1803)
islandica subsp. *crispiformis* (Räsänen) Kärnef. (1979)
* *juniperina* (L.) Ach. (1803)
nivalis (L.) Ach. (1803)
pinastri (Scop.) Gray (1821)
sepincola (Ehrh.) Ach. (1803)

Cetrelia Culb. & C. Culb. (1968)
olivetorum (Nyl.) Culb. & C. Culb. (1968)
C. cetrariooides (Del. ex Duby) Culb. & C. Culb. (1968)
C. chichitae Culb. & C. Culb. (1968)
C. monochorum (Zahlbr.) Culb. & C. Culb. (1968)

Chaenotheca (Th. Fr.) Th. Fr. (1860)
brachypoda (Ach.) Tibell (1987)
C. sulphurea (Retz) Middelb. & Mattsson (1987)
Coniocybe sulphurea (Retz) Nyl. (1886)
brunneola (Ach.) Müll. Arg. (1862)
chlorella (Ach.) Müll. Arg. (1862)
C. carthusiae (Harm.) Lettau (1912)
chrysoccephala (Turner ex Ach.) Th. Fr. (1860)
ferruginea (Turner ex Ach.) Mig. (1930)
furfuracea (L.) Tibell (1984)
Coniocybe furfuracea (L.) Ach. (1816)
gracilenta (Ach.) Mattsson & Middelb. (1987)
Coniocybe gracilenta (Ach.) Ach. (1816)
Cybebe gracilenta (Ach.) Tibell (1984)
hispidula (Ach.) Zahlbr. (1922)
laevigata Nádv. (1934)
phaeocephala (Turner) Th. Fr. (1860)
stemonea (Ach.) Müll. Arg. (1862)

trichialis (Ach.) Th. Fr. (1860)
xyloxena Nádv. (1934)

Chaenothecopsis Vainio (1927) [\pm L]
caespitosa (Phillips) D. Hawksw. (1980) [F]
* *debilis* (Turner & Borrer ex Sm.) Tibell (1975) [F]
epithallina Tibell (1975) [LF]
nigra Tibell (1987)
parasitaster (Bagl. & Car.) D. Hawksw. (1978) [LF]
pusilla (Ach.) A. Schmidt (1970) [F]
C. subpusilla (Vainio) Tibell (1975)
pusiola Ach. (Vainio (1927))
C. lignicola (Nádv.) A. Schmidt (1970)
? *rubescens* Vainio (1927)
savonica (Räsänen) Tibell (1984)
vainioana (Nádv.) Tibell (1979)
viridialba (Krempehl.) A. Schmidt (1970) [F]
viridireagens (Nádv.) A. Schmidt (1970) [LF]

Chiodecton Ach. (1814)
myrticola Fée (1825)
? *petraeum* Delise ex Duby (1830)

Chromatochlamys Trev. (1860)
larbalestieri (A. L. Sm.) Mayrh. & Poelt (1985)
Microglaena larbalestieri A.L. Sm. (1911)
muscorum (Fr.) Mayrh. & Poelt (1985)
Microglaena muscorum (Fr.) Th. Fr. (1860)
muscorum var. *octospora* (Nyl.) Mayrh. & Poelt (1985)

Chrysotrichia Mont. (1852)
candelaris (L.) Laundon (1981)
chlorina (Ach.) Laundon (1981)
chrysophthalma (P. James) P. James & Laundon (1981)

Cladina (Nyl.) Nyl. (1866) - see *Cladonia*.

Cladonia Hill ex Browne (1756)
arbuscula (Wallr.) Flotow (1839)
C. arbuscula subsp. *squarrosa* (Wallr.) Ruoss (1987)
azorica Ahti (1961)
bellidiflora (Ach.) Schaeerer (1823)
borealis Stenr. (1989)
botrytes (Hagen) Willd. (1728)
caespiticia (Pers.) Flörke (1828)
cariosa (Ach.) Sprengel (1827)
carneola (Fr.) Fr. (1831)
cenotea (Ach.) Schaeerer (1823)

- cervicornis* (Ach.) Flotow (1849)
· *C. cervicornis* subsp. *pulvinata* (Sandst.) Ahti (1983)
cervicornis subsp. *verticillata* (Hoffm.) Ahti (1980)
chlorophaea (Flörke ex Sommerf.) Sprengel (1827)
· *C. asahiniae* J.W. Thomson (1976)
· *C. cryptochlorophaea* Asah. (1940)
· *C. grayi* G.K. Merrill ex Sandst. (1929)
· *C. merochlorophaea* Asah. (1940)
· *C. merochlorophaea* var. *novochlorophaea* Sipman (1973)
ciliata Stirton (1888)
ciliata var. *tenuis* (Flörke) Ahti (1977)
coccifera (L.) Willd. (1787)
coniocraea (Flörke) Sprengel (1827)
conista A.W. Evans (1930)
convoluta (Lam.) Cout. (1913)
cornuta (L.) Hoffm. (1791)
crispata (Delise) Vainio (1887)
crispata var. *cetrariiformis* (Delise) Vainio (1887)
cyathomorpha Stirton ex W. Watson (1935)
deformis (L.) Hoffm. (1796)
digitata (L.) Hoffm. (1796)
diversa Asperges (1983)
fimbriata (L.) Fr. (1831)
firma (Nyl.) Nyl. (1861)
floerkeana (Fr.) Flörke (1828)
foliacea (Huds.) Willd. (1787)
fragilissima Østh. & P. James (1977)
furcata (Huds.) Schrader (1794)
glauca Flörke (1828)
gracilis (L.) Willd. (1787)
humilis (With.) Laundon (1984)
· *C. conoidea* Ahti (1980)
incrassata Flörke (1826)
luteoalba Wheldon & A. Wilson (1907)
macilenta Hoffm. (1796)
· *C. bacillaris* (Leighton) Arnold (1880)
macrophylla (Schaerer) Stenhammar (1865)
maxima (Asah.) Ahti (1978)
mediterranea Duvign. & des Abb. (1947)
metacorallifera Asah. (1939)
mitis Sandst. (1918)
· *Cladina mitis* (Sandst.) Hustich (1951)
? *C. arbuscula* subsp. *mitis* (Sandst.) Ruoss (1987)
norvegica Tønsb. & Holien (1984)
ochrochlora Flörke (1828)
parasitica (Hoffm.) Hoffm. (1796)

- peziziformis (With.) Laundon (1984)
C. capitata (Michaux) Sprengel (1827)
phylophora Hoffm. (1796)
pleurota (Flörke) Schaerer (1850)
pocillum (Ach.) O.J. Rich (1878)
C. pyxidata subsp. *pocillum* (Ach.) Dahl (1950)
polydactyla (Flörke) Sprengel (1827)
polydactyla var. *umbricola* (Tønsb. & Ahti) Coppins (1993)
C. umbricola Tønsb. & Ahti (1980)
portentosa (Dufour) Coem. (1865)
C. portentosa f. *subimpexa* (Duvign.) Ahti (1978)
pyxidata (L.) Hoffm. (1796)
ramulosa (With.) Laundon (1984)
C. anomaea (Ach.) Ahti & P. James (1980)
rangiferina (L.) Weber ex Wigg. (1780)
rangiformis Hoffm. (1796)
rei Schaerer (1823)
scabriuscula (Delise) Nyl. (1876)
squamosa (Scop.) Hoffm. (1796)
squamosa var. *subsquamosa* (Nyl. & Leighton) Vainio (1881)
* *stellaris* (Opiz) Pouzar & Vězda (1971)
stereoclada des Abb. (1946)
strepsilis (Ach.) Vainio (1894)
stricta (Nyl.) Nyl. (1869)
stygia (Fr.) Ruoss (1985)
subcervicornis (Vainio) Kernst. (1900)
subrangiformis Sandst. (1924)
C. furcata subsp. *subrangiformis* (Scriba ex Sandst.) Pišút (1961)
subulata (L.) Weber ex Wigg. (1780)
sulphurina (Michaux) Fr. (1831)
sympycarpa (Ach.) Fr. (1826)
uncialis (L.) Weber ex Wigg. (1780)
uncialis subsp. *biuncialis* (Hoffm.) M. Choisy (1951)
zoppii Vainio (1920)

Literature: Stenroos (1989a, 1989b).

Clathroporina Müll. Arg. (1882) - see *Belonia*.

Clauroxia D. Hawksw. (1988)
chalybeoides (Nyl.) D. Hawksw. (1988)
Lecidea chalybeoides Nyl. (1875)
Pseudolecidea chalybeoides (Nyl.) Clauz. & Roux (1984)

Clauzadea Hafellner & Bellem. (1984)
immersa (Hoffm.) Hafellner & Bellem. (1984)
Lecidea immersa (Hoffm.) Ach. (1803)

metzleri (Körber) Clauz. & Roux ex D. Hawksw. (1992)

Lecidea metzleri (Körber) Th. Fr. (1874)

monticola (Ach.) Hafellner & Bellem. (1984)

Lecidea monticola (Ach.) Schaeerer (1833)

Clauzadeana Roux (1984)

macula (Taylor) Coppins & Rambold (1989)

Aspicilia morioides Blomb. ex Arnold (1881)

Clauzadeana instratula (Nyl.) Roux (1984)

Lecidea instratula Nyl. (1878)

Lecidea macula Taylor (1836)

Lecidea pissodes Stirton (1877)

Cliostomum Fr. (1825)

corrugatum (Ach.) Fr. (1831)

C. graniforme (Hagen) Coppins (1980)

griffithii (Sm.) Coppins (1980)

Coccotrema Müll. Arg. (1888)

citrinescens P. James & Coppins (1992)

Coelocaulon Link (1833)

aculeatum (Schreber) Link (1833)

muricatum (Ach.) Laundon (1984)

Collema Weber ex Wigg. (1780)

auriforme (With.) Coppins & Laundon (1984)

C. auriculatum Hoffm. (1796)

bachmanianum (Fink) Degel. (1954)

callopismum Massal. (1856)

callopismum var. *rhyparodes* (Nyl.) Degel. (1954)

ceraniscum Nyl. (1865)

confertum Arnold (1867)

* *conglomeratum* Hoffm. (1796)

crispum (Huds.) Weber ex Wigg. (1780)

crispum var. *metzleri* (Arnold) Degel. (1954)

cristatum (L.) Weber ex Wigg. (1780)

cristatum var. *marginale* (Huds.) Degel. (1954)

dichotomum (With.) Coppins & Laundon (1984)

fluviale (Huds.) Steudel (1824)

fasciculare (L.) Weber ex Wigg. (1780)

flaccidum (Ach.) Ach. (1810)

fragile Taylor (1836)

fragrans (Sm.) Ach. (1814)

furfuraceum (Arnold) Du Rietz (1929)

fuscovirens (With.) Laundon (1984)

C. tuniforme (Ach.) Ach. (1810)

glebulentum (Nyl. ex. Crombie) Degel. (1952)

latzelii Zahlbr. (1909)
limosum (Ach.) Ach. (1810)
multipartitum Sm. (1814)
nigrescens (Huds.) DC. (1805)
occultatum Bagl. (1861)
parvum Degel. (1954)
polycarpon Hoffm. (1796)
subflaccidum Degel. (1974)
subnigrescens Degel. (1954)
tenax (Swartz) Ach. (1810)
tenax var. *ceranoides* (Borrer) Degel. (1954)
tenax var. *corallinum* (Massal.) Degel. (1954)
tenax var. *vulgare* (Schreber) Degel. (1954)
undulatum Lauter ex Flotow. (1850)
undulatum var. *granulosum* Degel. (1954)

Conioocybe Ach. (1816)

C. furfuracea (L.) Ach. (1816) = *Chaenotheca*
C. gracilenta Ach. (1816) = *Chaenotheca*
C. pallida (Pers.) Fr. (1824) = *Sclerophora*
C. peronella (Ach.) Tibell (1978) = *Sclerophora*.

Coriscium Vainio (1890) - see *Omphalina*.

Cornicularia (Schreber) Hoffm. (1794)
normoerica (Gunn.) Du Rietz (1926)

Cryptolechia Massal. (1853)
carneolutea (Turner) Massal. (1853)

Cryptothelae Th. Fr. (1860)
 * *rhodosticta* (Taylor) Henssen (1990)

Cyphelium Ach. (1815)
inquinans (Sm.) Trevisan (1862)
marcianum B. de Lesd. (1908)
notarisii (Tul.) Blomb. & Forss. (1880)
sessile (Pers.) Trevisan (1862)
tigillare (Ach.) Ach. (1815)

Cystocoleus Thwaites (1849)
ebeneus (Dillwyn) Thwaites (1849)
C. niger (Huds.) Hariot (1890)

Degelia Arvidsson & D. Galloway (1981)
atlantica (Degel.) P.M. Jørg. & P. James (1990)
Parmeliella atlantica Degel. (1935)
ligulata P.M. Jørg & P. James (1990)

plumbea (Lightf.) P.M. Jørg. & P. James (1990)
Parmeliella plumbea (Lightf.) Vainio (1890)

Dendriscocaulon Nyl. (1888) - see *Lobaria*.

Dermatocarpon Eschw (1924)

arnoldianum Degel. (1934)
intestiniforme (Körber) Hasse (1912)
? leptophyloides (Nyl.) Zahlbr. (1922)
leptophyllum (Ach.) G. Lång (1912)
luridum (With.) Laundon (1984)
D. weberi (Ach.) Mann (1825)
meiophyllizum Vainio (1921)
miniatum (L.) Mann (1825)
miniatum var. *complicatum* (Lightf.) Hellbom (1870)
rivulorum (Arnold) Dalla Torre & Sarnth. (1902)

D. hepaticum auct., non (Ach.) Th. Fr. (1860) = *Catapyrenium*.

Dictyonema C. Ag. (1821)

interruptum (Carm. ex Hook.) Parm. (1978)

Dimerella Trevisan (1880)

lutea (Dickson) Trevisan (1880)
pineti (Ach.) Vězda (1975)
D. diluta (Pers.) Trevisan (1880)

Diploicia Massal. (1852)

canescens (Dickson) Massal. (1852)

Diploschistes Norman (1853)

caesioplumbeus (Nyl.) Vainio (1921)
gypsaceus (Ach.) Zahlbr. (1892)
muscorum (Scop.) R. Sant. (1980)
D. scruposus var. *bryophilous* (Ehrh. ex Ach.) Müll. Arg. (1892)
scruposus (Schreber) Norman (1853)

Diplotomma Flotow (1849)

alboastrum (Hoffm.) Flotow (1849)
Buellia geophila auct. brit., non (Flörke ex Sommerf.) Lynge (1937)
chlorophaeum (Hepp ex Leighton) Szat. (1956)
epipodium (Ach.) Arnold (1869)
muronorum (Massal.) Coppins (1980)

Dirina Fr. (1825)

massiliensis Durieu & Mont. (1847)
D. repanda (Fr.) Nyl. (1836)

massiliensis f. *sorediata* (Müll. Arg.) Tehler (1983)

D. repanda f. *stenhammarii* (Fr. ex Stenhammar) Clauz. & Roux (1975)

DiscosIELla H. Sydow & Sydow (1912) - see *Strigula*.

Eiglera Hafellner (1984)

flavida (Hepp) Hafellner (1984)

Aspicilia flavida (Hepp) Rehm (1863)

Encephalographa Massal. (1854) - see *Poeltinula*.

E. interjecta (Leighton) Th. Fr. (1874) is included in *Melaspilea*.

Endocarpon Hedwig (1789)

adscendens (Anzi) Müll. Arg. (1881)

* *pusillum* Hedwig (1789)

* *pusillum* var. *pallidum* (Ach.) Körber (1855)

Enterographa Fée (1824)

crassa (DC.) Fée (1824)

elaborata (Lyell ex Leighton) Coppins & P. James (1979)

hutchinsiae (Leighton) Massal. (1860)

sorediata Coppins & P. James (1979)

zonata (Körber) Källsten (1989)

Opegrapha zonata Körber (1855)

Eopyrenula R. Harris (1973)

avellanae Coppins (1992)

grandicula Coppins (1992)

leucoplaca (Wallr.) R. Harris (1973)

septemseptata Coppins (1992)

Ephebe Fr. (1825)

hispidula (Ach.) Horwood (1913)

lanata (L.) Vainio (1888)

Epigloea Zukal (1890)

grummanni Döbbeler (1984)

E. bactrospora auct. brit., non Zukal (1889)

medioincrassata (Grumm.) Döbbeler (1984)

Vorarlbergia medioincrassata Grumm. (1969)

soleiformis Döbbeler (1984)

Epilichen Clem. (1909)

scabrosus (Ach.) Clem. (1909)

Euopsis Nyl. (1875)

granatina (Sommerf.) Nyl. (1875)

Pyrenopsis granatina (Sommerf.) Nyl. (1866)

pulvinata (Schaerer) Vainio (1881)

Pyrenopsis pulvinata (Schaerer) Th. Fr. (1866)

Evernia Ach. (1810)

prunastri (L.) Ach. (1810)

E. prunastri f. *herinii* (Duvign.) D. Hawksw. (1980)

Farnoldia Hertel (1983)

jurana (Schaerer) Hertel (1983)

Melanolecia jurana (Schaerer) Hertel (1981)

Tremolecia jurana (Schaerer) Hertel (1977)

Fellhanera Vězda (1986)

bouteillei (Desm.) Vězda (1986)

Catillaria bouteillei (Desm.) Zahlbr. (1902)

subtilis (Vězda) Diederich & Sérusiaux (1991)

Bacidia subtilis Vězda (1961)

F. vezdae (Coppins & P. James) V. Wirth (1987) = *Bacidia*.

Flavoparmelia Hale (1986) - see *Parmelia*.

Foraminella Fricke Meyer (1982) - see *Parmeliopsis*.

Fulglesia Massal. & de Not. (1855)

bracteata (Hoffm.) Räsänen (1931) var. *alpina* (Th. Fr.) Räsänen (1943)

fulgens (Swartz) Elenkin (1907)

Fuscidea V. Wirth & Vězda (1972)

arboricola Coppins & Tønsb. (1992)

austera (Nyl.) P. James (1980)

? *F. taeniarum* (Malme) V. Wirth & Vězda (1972)

cyathoides (Ach.) V. Wirth & Vězda (1972)

F. cyathoides var. *corticola* (Fr.) Kalb (1976)

cyathoides var. *sorediata* (Magnusson) Poelt (1978)

gothoburgensis (Magnusson) V. Wirth & Vězda (1972)

F. gothoburgensis f. *maculosa* (Magnusson) Poelt (1981)

hibernica P. James & Poelt (1992)

intercincta (Nyl.) Poelt (1978)

F. atlantica (Magnusson) P. James & Poelt (1981)

kochiana (Hepp) V. Wirth & Vězda (1972)

lightfootii (Sm.) Coppins & P. James (1978)

lugubris (Sommerf.) P. James & Purvis (1992)

Ropalospora lugubris (Sommerf.) Poelt (1980)

lygaea (Ach.) V. Wirth & Vězda (1972)

F. periplaca (Nyl.) V. Wirth & Vězda (1972)

F. tenebrica (Nyl.) V. Wirth & Vězda (1972)

mollis (Wahlenb.) V. Wirth & Vězda (1972)

praeruptorum (Du Rietz & Magnusson) V. Wirth & Vězda (1972)

pusilla Tønsb. (1992)

recensa (Stirton) Hertel, V. Wirth & Vězda (1972)

viridis Tønsb. (1984)

Ropalospora viridis (Tønsb.) Tønsb. (1992)

Geltingia Alstrup & D. Hawksw. (1990) [LF]

associata (Th. Fr.) Alstrup & D. Hawksw. (1990) [LF]

Lecidea associata Th. Fr. (1867)

groenlandiae Alstrup & D. Hawksw. (1990) [LF]

Gerronema Singer (1951) - see *Omphalina*.

Glaucomaria M. Choisy (1929) - see *Lecanora*.

Gomphillus Nyl. (1853)

calycoides (Del.) Nyl. (1853)

Graphina Müll. Arg. (1880)

anguina (Mont.) Müll. Arg. (1882)

pauciloculata Coppins & P. James (1978)

ruiziana (Fée) Müll. Arg. (1887)

Graphis Adans. (1763)

alboscripta Coppins & P. James (1992)

elegans (Borrer ex Sm.) Ach. (1814)

scripta (L.) Ach. (1809)

Gyalecta Ach. (1808)

biformis (Körber) H. Olivier (1911)

derivata (Nyl.) H. Olivier (1911)

G. truncigena var. *derivata* (Nyl.) Boistel (1903)

flotowii Körber (1855)

foveolaris (Ach.) Schaeerer (1836)

geoica (Wahlenb.) Ach. (1808)

jenensis (Batsch) Zahlbr. (1924)

jenensis var. *macrospora* Vězda (1969)

truncigena (Ach.) Hepp (1853)

ulmi (Swartz) Zahlbr. (1905)

Gyalidea Lettau (1937)

diaphana (Körber ex Nyl.) Vězda (1978)

fritzei (Stein) Vězda (1966)

hyalinescens (Nyl.) Vězda (1966)

lecidopsis (Massal.) Lettau ex Vězda (1966)

roseola (Arnold) Lettau ex Vězda (1966)

subscutellaris (Vězda) Vězda (1966)

Gyalideopsis Vězda (1972)

anastomosans P. James & Vězda (1972)

muscicola P. James & Vězda (1972)

scotica P. James (1975)

Haematomma Massal. (1852)

leprarioides (Vainio) Vainio (1934)

ochroleucum (Necker) Laundon (1970)

ochroleucum var. porphyrium (Pers.) Laundon (1970)

H. caesium Coppins & P. James (1978) = *Mycoblastus*.*H. elatinum* (Ach.) Massal. (1854) = *Loxospora**H. lapponica* Räsänen (1931) = *Ophioparma**H. ventosum* (L.) Massal. (1852) = *Ophioparma*.***Hafellia* Kalb, Mayrh. & Scheid. (1986) - see *Buellia*.****Halecania M. Mayrh. (1987)***alpivaga* (Th. Fr.) M. Mayrh. (1987)*Lecania alpivaga* Th. Fr. (1871)*ralfsii* (Salwey) M. Mayrh. (1987)*Catillaria jejuna* (Nyl.) P. James (1965)*Lecania ralfsii* (Salwey) A.L. Sm. (1918)*Lecidea muddii* Salwey ex Mudd. (1853)*rhypodiza* (Nyl.) Coppins (1989)*Catillaria rhypodiza* (Nyl.) Zahlbr. (1926)*spodomela* (Nyl.) M. Mayrh. (1987)*Lecania spodomela* (Nyl.) A.L. Sm. (1918)*Lecidea nigricans* Nyl. (1876)*viridescens* Coppins & P. James (1989)***Herteliana* P. James (1980)***taylorii* (Salwey) P. James (1980)***Heterodermia* Trevisan (1868)***isidiophora* (Vainio) Awasthi (1973)*leucomelos* (L.) Poelt (1965)*obscurata* (Nyl.) Trevisan (1869)*propagulifera* (Vainio) Day (1977)***Hulia* Zahlbr. (1930) - see *Porpidia*.***H. consentiens* (Nyl.) Hertel (1980) is included in *Amygdalaria*.***Hymenelia* Krempelh. (1852)***lacustris* (With.) M. Choisy (1949)*prevostii* (Duby) Krempelh. (1852)

Hyperphyscia Müll. Arg. (1894)

adglutinata (Flörke) Mayrh. & Poelt (1979)

Physciopsis adglutinata (Flörke) M. Choisy (1950)**Hypocenomyce** M. Choisy (1951)

anthracophila (Nyl.) P. James & G. Schneider (1980)

caradocensis (Leighton ex Nyl.) P. James & G. Schneider (1980)

friesii (Ach.) P. James & G. Schneider (1980)

leucococca R. Sant. (1986)

scalaris (Ach. ex Lilj.) M. Choisy (1951)

soropora (Vainio) P. James & Poelt (1981)

xanthococca (Sommerf.) P. James & G. Schneider (1980)

Hypogymnia (Nyl.) Nyl. (1896)

bitteriana (Zahlbr.) Räsänen (1947)

intestiniformis (Vill.) Räsänen (1943)

physodes (L.) Nyl. (1896)

tubulosa (Schaerer) Havaas (1918)

Hypotrachyna (Vainio) Hale (1974) - see *Parmelia*.**Icmadophila** Trevisan (1852)

ericetorum (L.) Zahlbr. (1895)

Immersaria Rambold & Pietschmann (1989)

athroocarpa (Ach.) Rambold & Pietschm. (1989)

Lecidea athroocarpa (Ach.) Ach. (1803)*Porpidia athroocarpa* (Ach.) Hertel & Rambold (1985)**Imshaugia** S. Meyer (1985)

aleurites (Ach.) S. Meyer (1985)

Parmeliopsis aleurites (Ach.) Nyl. (1866)**Ionaspis** Th. Fr. (1871)

cyanocarpa (Anzi) Jatta (1910)

epulotica (Ach.) Arnold (1871)

I. epulotica var. *crustosa* Magnusson (1933)*I. rhodopsis* (Sommerf.) Blomb. & Forss. (1880)

heteromorpha (Krempelh.) Arnold (1874)

melanocarpa (Krempelh.) Arnold (1887)

odora (Ach.) B. Stein (1879)

suaveolens (Schaerer) Arnold (1874)

Japewia Tønsb. (1990)

carrollii (Coppins & P. James) Tønsb. (1990)

Lecidea carrollii Coppins & P. James (1979)*Lecidea rufofusca* auct. brit., non (Anzi) Nyl. (1869)*Lecidea tenebricosa* auct. brit., non (Ach.) Nyl. (1861)

subaurifera Muhr & Tønsb. (1990)
tornoensis (Nyl.) Tønsb. (1990)
Lecidea tornoensis Nyl. (1859)
Mycoblastus tornoensis (Nyl.) R. Anderson (1974)

Julella Fabre (1879)
sericea (Massal.) Coppins (1992)
Polyblastiopsis sericea (Massal.) Zahlbr. (1903)

Kiliasia Hafellner (1984) - see *Catillaria*.

Koerberiella B. Stein (1879)
wimmeriana (Körber) B. Stein (1879)
Aspicilia leucophyma (Leighton) Hue (1912)

Lasallia Mérat (1821)
pustulata (L.) Mérat (1821)

Lauderlindsaya J.C. David & D. Hawksw. (1989) (\pm L]
borreri (L.R. Tulasne) J.C. David & D. Hawksw. (1989) [LF]
chlorococca (Leighton) Diederich & Sérusiaux (1993)
L. erichsenii (Keissler) Diederich & Sérusiaux (1991)
Normandina erichsenii (Keissler) Aptroot (1991)
Sphaerulina chlorococca (Leighton) R. Sant. (1976)

Literature: Diederich & Sérusiaux (1993)

Lecanactis Fr. (1825)
abietina (Ach.) Körber (1855)
abscondita (Th. Fr.) Lojka (1874)
amylacea (Ehrh. ex Pers.) Arnold (1880)
dilleniana (Ach.) Körber (1855)
grumulosa (Dufour) Fr. (1831)
L. delimitis (Nyl.) A.L. Sm. (1911)
L. mirifica (Stirton) A.L. Sm. (1926)
L. monstrosa Bagl. (1857)
L. nothiza (Nyl.) P. James (1980)
hemisphaerica Laundon (1976)
latebrarum (Ach.) Arnold (1885)
lyncea (Sm.) Fr. (1831)
Opegrapha lyncea (Sm.) Borrer ex Hook. (1833)
premnea (Ach.) Arnold (1861)
L. plocina auct., non (Ach.) Massal (1860)
subabietina Coppins & P. James (1979)

L. umbrina Coppins & P. James (1979) = *Schismatomma*
L. homalotropa (Nyl.) Gilenstam (1969) = *Bactrospora*.

Lecania Massal. (1853)*aipospila* (Wahlenb.) Th. Fr. (1867)*atrynoides* Knowles (1913)*baeomma* (Nyl.) P. James & Laundon (1980)*chlorotiza* (Nyl.) P. James (1992)*Catillaria chlorotiza* (Nyl.) P. James (1965)* *coeruleorubella* (Mudd) M. Mayrh. (1988)*coerulescens* Mudd (1861)*cuprea* (Massal.) v.d. Boom & Coppins (1992)*Bacidia cuprea* (Massal.) Lettau (1912)*Catillaria umbraticula* (Nyl.) P. James (1965)*cyrtella* (Ach.) Th. Fr. (1871)*Lecania sambucina* (Körber) Arnold (1884)*cyrtellina* (Nyl.) Sandst. (1912)*Lecidea albohyalina* auct. brit., non (Nyl.) Th. Fr. (1875)*Lecidea meiocarpa* auct. brit., non Nyl. (1876)*erysibe* (Ach.) Mudd (1861)*Lecania erysibe* f. *sorediata* Laundon (1967)* *fuscella* (Schaerer) Körber (1855)*hutchinsiae* (Nyl.) A. L. Sm. (1918)*Catillaria littorella* auct. brit., non (Nyl.) Zahlbr. (1926)*inundata* (Hepp ex Körber) M. Mayrh. (1987)*nylanderiana* Massal. (1856)* *olivacella* (Nyl.) Zahlbr. (1928)*rabenhorstii* (Hepp) Arnold (1848)*suavis* (Müll. Arg.) Mig. (1926)*sylvestris* (Arnold) Arnold (1884)*turicensis* (Hepp) Müll. Arg. (1862)*Lecania albariella* (Nyl.) Müll. Arg. (1880)*L. alpivaga* Th. Fr. (1871) = *Halecania**L. helicopis* (Wahlenb.) Clauz. & Roux (1985) = *Lecanora**L. ralfsii* (Salwey) A.L. Sm. (1918) = *Halecania**L. spodomela* (Nyl.) A.L. Sm. (1918) = *Halecania**L. actea* (Nyl.) B. de Lesd. (1910) = *Catillaria*.**Lecanora** Ach. (1810)*achariana* A.L. Sm. (1918)*actophila* Wedd. (1875)*agardhiana* Ach. (1814)*aitema* (Ach.) Hepp (1853)? *aitema* var. *saepincola* (Ach.) Hedl. (1892)*albescens* (Hoffm.) Branth & Rostrup (1869)*alboflavida* Taylor (1836)*andrewii* B. de Lesd. (1913)*argentata* (Ach.) Malme (1897)*L. subfuscata* Magnusson (1932)*atrosulphurea* (Wahlenb.) Ach. (1814)

- cadubriae (Massal.) Hedl. (1892)
Lecidea cadubriae (Massal.) Th. Fr. (1874).
caesiosora Poelt (1966)
campestris (Schaerer) Hue (1888)
campestris subsp. dolomitica O.L. Gilbert (1984)
carpinea (L.) Vainio (1888)
cenisia Ach. (1810)
cenisia var. soredians Suza (1928)
chlarotera Nyl. (1872)
chlorophaeodes Nyl. (1873)
cinereofusca Magnusson (1932)
conferta (Duby ex Fr.) Grognot (1863)
confusa Almb. (1955)
* congesta Clauz. & Vézda (1969)
conizaeoides Nyl. ex Crombie (1885)
crenulata Hook. (1833)
dispersa (L.) Sommerf. (1826)
epanora (Ach.) Ach. (1810)
epibryon (Ach.) Ach. (1810)
expallens Ach. (1810)
farinaria Borrer (1834)
frustulosa (Dickson) Ach. (1810)
fugiens Nyl. (1873)
* fuscescens (Sommerf.) Nyl. (1871)
Lecidea fuscescens Sommerf. (1824)
gangaleoides Nyl. (1872)
gisleriana Müll. Arg. (1874)
L. gisleri Poelt & Ullrich (1964)
handelii Steiner (1909)
helicopis (Wahlenb.) Ach. (1814)
Lecania helicopis (Wahlenb.) Clauz. & Roux (1985)
horiza (Ach.) Lindsay (1869)
Lecanora parisiensis Nyl. (1866)
intricata (Ach.) Ach. (1810)
intumescens (Rebent.) Rabenh. (1845)
jamesii Laundon (1963)
laevis Poelt (1952)
leptacina Sommerf. (1826)
marginata (Schaerer) Hertel & Rambold (1985)
Lecidea marginata Schaerer (1818)
mughicola Nyl. (1872)
muralis (Schreber) Rabenh. (1845)
orosthea (Ach.) Ach. (1810)
Lecidea orosthea (Ach.) Ach. (1803)
pallida (Schreber) Rabenh. (1845), non Chevall. (1826)
Lecanora albella (Pers.) Ach. (1810)
persimilis (Th. Fr.) Nyl. (1876)
piniperda Körber (1859)

- poliophaea (Wahlenb.) Ach. (1810)
- * populincola (DC.) Duby (1830)
- polytropa (Hoffm.) Rabenh. (1845)
- praepostera Nyl. (1873)
 - ? *L. schistina* (Nyl.) Nyl. ex Crombie (1894)
- * pruinosa Chaub. (1821)
- pulicaris (Pers.) Ach. (1814)
- quercicola Coppins & P. James (1979)
- rugosella Zahlbr. (1928)
- rupicola (L.) Zahlbr. (1928)
 - Aspicilia lindsayi* W. Watson (1939)
- rupicola var. efflorens Leuckert & Poelt (1989)
 - L. rupicola* f. *sorediata* (Flotow) Zahlbr. (1928)
- saligna (Schrader) Zahlbr. (1928)
- salina Magnusson (1926)
- sambuci (Pers.) Nyl. (1861)
- sarcopisoides (Massal.) A.L. Sm. (1918)
- soralifera (Suza) Räsänen (1931)
- stenotropa Nyl. (1872)
- straminea Wahlenb. ex Ach. (1810)
- strobilina (Sprengel) Kieffer (1895)
- subaurea Zahlbr. (1928)
- subcarnea (Lilj.) Ach. (1810)
- subintricata (Nyl.) Th. Fr. (1871)
- sublivescens (Nyl. ex Crombie) A.L. Sm. (1918)
 - Lecidea sublivescens* (Nyl. ex Crombie) P. James (1965)
- sulphurea (Hoffm.) Ach. (1810)
 - Lecidea sulphurea* (Hoffm.) Wahlenb. (1812)
- swartzii (Ach.) Ach. (1810)
 - L. radiosa* var. *karellica* Räsänen (1934)
 - L. subradiosa* Nyl. (1872)
- symmicta (Ach.) Ach. (1814)
- tenera (Nyl.) Crombie (1874)
- varia (Hoffm.) Ach. (1810)

- L. atra* Huds. Ach. (1810) = *Tephromela*
- L. atriseda* (Fr.) Nyl. (1857) = *Protoparmelia*
- L. badia* (Hoffm.) Ach. (1810) = *Protoparmelia*
- L. contracta* (Th. Fr.) Zahlbr. (1928) = *Rimularia*
- L. curvescens* (Mudd) Nyl. (1865) = *Bryonora*
- L. grumosa* (Pers.) Du Rietz (1926) = *Tephromela*
- L. intercedens* Magnusson (1930) = *Rimularia*
- L. nitens* auct., non Ach. (1814) = *Protoparmelia*
- L. ochrococca* (Nyl.) Clauz. & Roux (1985) = *Protoparmelia*
- L. oleagina* Harm. (1913) = *Protoparmelia*
- L. picea* (Dickson) Nyl. (1986) = *Protoparmelia*
- L. psarophana* Nyl. (1872) = *Protoparmelia*.

Lecanoropsis M. Choisy (1949) - see *Lecanora*.

Lecidea Ach. (1803)

* *advertisens* Nyl. (1866)

ahlesii (Körber) Nyl. (1872)

L. valentior Nyl. (1877)

antilogia Stirton (1877)

auriculata Th. Fr. (1861)

berengeriana (Massal.) Nyl. (1866)

botryosa (Fr.) Th. Fr. (1874)

brachyspora (Th. Fr.) Nyl. (1887)

caesioatra Schaeer (1818)

? *cetrariicola* Lindsay (1869) [LF]

* *commaculans* Nyl. (1868)

confluens (Web.) Ach. (1803)

diducens Nyl. (1865)

L. auriculata var. *diducens* (Nyl.) Th. Fr. (1874)

doliiformis Coppins & P. James (1992)

endomelaena Leighton (1876)

erythrophaea Flörke (1826)

L. hyalinella (Körber) Jatta (1889)

exigua Chaub. (1821)

fuliginosa Taylor (1836)

fuscoatra (L.) Ach. (1803)

L. fuscoatra var. *grisella* (Flörke ex Schaeer) Nyl. (1856)

hypnorum Lib. (1853)

hypopta Ach. (1803)

inops Th. Fr. (1874)

insidiosa Th. Fr. (1868)

lactea Flörke ex Schaeer (1828)

L. demolita auct. brit., non Vainio (1934)

L. lapicida var. *lactea* (Flörke ex Schaeer) V. Wirth (1987)

lapicida (Ach.) Ach. (1803)

L. contiguela Nyl. (1873)

L. luteoatra Nyl. (1873)

L. subcinerascens auct. brit., non Nyl. (1877)

lichenicola (A. L. Sm. & Ramsb.) D. Hawksw. (1978)

? *L. obsoleta* Nyl. (1865)

limosa Ach. (1810)

L. stenotera auct. brit., non (Nyl.) Nyl. (1866)

lithophila (Ach.) Ach. (1814)

matildae Magnusson (1956)

nigrogrisea Nyl. (1879)

nylanderi (Anzi) Th. Fr. (1874)

* *paraclitica* Nyl. (1872)

paupercula Th. Fr. (1874)

pernigra Hertel (1975)

phaeops Nyl. (1858)

plana (Lahm) Nyl. (1872)
polycarpella Erichsen (1929)
pycnocarpa (Körber) Ohlert (1870)
L. segregans Nyl. (1866)
L. tabidula Nyl. (1867)
pycnocarpa f. *sorediata* Coppins & Fryday (1992)
sanguineoatra auct., non (Wulff) Ach. (1803)
sarcogynoides Körber (1855)
silacea (Ach.) Ach. (1803)
turgidula Fr. (1824)

L. aeruginosa Borrer (1831) = *Trapeliopsis*
L. aglaea Sommerf. (1826) = *Tephromela*
L. albohyalina auct. brit., non (Nyl.) Th. Fr. (1875) = *Lecania*
L. aphana Nyl. (1867) = *Catillaria*
L. aphanoides Nyl. (1868) = *Micarea*
L. armeniaca (DC.) Fr. (1825) = *Tephromela*
L. assimilata Nyl. (1861) = *Micarea*
L. associata Th. Fr. (1867) = *Geltingia*
L. athroocarpa (Ach.) Ach. (1803) = *Immersaria*
L. atrofulva Sommerf. (1826) = *Miriquidica*
L. botryiza Nyl. ex Stirton (1873) = *Micarea*
L. cadubriae (Massal.) Th. Fr. (1874) = *Lecanora*
L. carrollii Coppins & P. James (1979) = *Japewia*
L. chalybeioides Nyl. (1875) = *Clauroxia*
L. cinnabarina auct. brit. p.p., non Sommerf. (1823) = *Pertusaria*
L. confusior Nyl. (1874) = *Catillaria*
L. cuprea Sommerf. (1826) = *Biatora*
L. dolosa (Nyl.) Vainio (1983) = *Lecidella*
L. efflorescens (Hedl.) Erichsen (1932) = *Biatora*
L. epixanthoidiza auct., non Nyl. (1875) = *Biatora*
L. erratica Körber (1861) = *Micarea*
L. furvella Nyl. ex Mudd (1861) = *Rimularia*
L. fuscescens Sommerf. (1824) = *Lecanora*
L. fuscocinerea auct., non Nyl. (1852) = *Rimularia*
L. fuscocinerea Nyl. (1852), non auct. brit. = *Schaereria*
L. gelatinosa Flörke (1809) = *Trapeliopsis*
L. granulosa (Hoffm.) Ach. (1803) = *Trapeliopsis*
L. griseoatra (Flotow) Schaeerer (1850) = *Miriquidica*
L. grunosa Leighton (1876) = *Agyrium*
L. gyrisans Nyl. (1861) = *Rimularia*
L. gyrophorica Tønsb. (1992) = *Biatora*
L. humigena Taylor (1847) = *Aphanopsis*
L. illita Nyl. (1879) = *Rimularia*
L. immersa (Hoffm.) Ach. (1803) = *Clauzadea*
L. inquinans (Tul.) Nyl. (1856) = *Micarea*
L. insularis Nyl. (1852) = *Rimularia*
L. instratula Nyl. (1878) = *Clauzadeana*

- L. leucophaea* (Flörke ex Rabenh.) Nyl. (1870) = *Miriquidica*
L. limborina (Nyl.) Lamy (1878) = *Rimularia*
L. lulensis Hellbom (1866) = *Miriquidica*
L. macula Taylor (1836) = *Clauzadeana*
L. marginata Schaerer (1818) = *Lecanora*
L. meiocarpa auct. brit., non Nyl. (1876) = *Lecania*
L. melaphana Nyl. (1869) = *Micarea*
L. metzleri (Körber) Th. Fr. (1874) = *Clauzadea*
L. modesta Müll. Arg. (1871) = *Catillaria*
L. monticola (Ach.) Schaerer (1833) = *Clauzadea*
L. muddii Salwey ex Mudd. (1853) = *Halecania*
L. mullensis Stirton (1877) = *Rimularia*
L. nigrificans Nyl. (1876) = *Halecania*
L. nigroglomerata Leighton ex Crombie (1871) = *Lecidella*
L. nigroleprosa (Vainio) Magnusson (1927) = *Miriquidica*
L. ochrococca Nyl. (1860) = *Protoparmelia*
L. oligotropha Laundon (1960) = *Placynthiella*
L. orosthea (Ach.) Ach. (1803) = *Lecanora*
L. pedatula Nyl. (1876) = *Lepraria*
L. pilati (Hepp) Körber (1861) = *Adelotelia*
L. pisoides Stirton (1877) = *Farnoldia*
L. punctum (Massal.) Jatta (1900) = *Phaeopyxis*
L. rhizobola Nyl. (1865) = *Acarospora*
L. rufofusca auct. brit., non (Anzi) Nyl. (1869) = *Japewia*
L. rusticella Nyl. (1878) = *Micarea*
L. scotinodes Nyl. (1873) = *Catillaria*
L. speirea (Ach.) Ach. (1803) = *Porpidia*
L. subgyratula Nyl. (1873) = *Rimularia*
L. sublivescens (Nyl. ex Crombie) P. James (1965) = *Lecanora*
L. submoestula Nyl. (1876) = *Micarea*
L. subplumbea Anzi (1868) = *Miriquidica*
L. sulphurea (Hoffm.) Wahlenb. (1812) = *Lecanora*
L. supersparsa Nyl. (1865) = *Carbonea*
L. tenebricosa auct. brit., non (Ach.) Nyl. (1861) = *Japewia*
L. tornoensis Nyl. (1859) = *Japewia*
L. uliginascens Stirton (1878) = *Placynthiella*
L. uliginosa (Schrader) Ach. (1803) = *Placynthiella*
L. umbonata auct. brit., non (Hepp) Mudd (1861) = *Cecidonia*
L. umbonella Nyl. (1866) = *Cecidonia*
L. umbonella var. *alumnula* (Nyl.) Hertel (1970) = *Cecidonia*
L. vernalis (L.) Ach. (1803) = *Biatora*
L. viridescens (Schrader) Ach. (1803) = *Trapeliopsis*
L. vitellinaria Nyl. (1852) = *Carbonea*
L. vorticosa (Flörke) Körber (1855) = *Carbonea*.

Lecidella Körber (1855)

- anomaloides (Massal.) Hertel & Kilias (1980)
- asema (Nyl.) Knoph & Hertel (1990)
 - L. subincongrua* (Nyl.) Hertel & Leuckert (1969)
 - L. effugiens* (B. Nils.) Knoph & Hertel (1990)
 - L. elaeochromoides* (Nyl.) Knoph & Hertel (1990)
 - L. polyantha* Taylor ex Leighton (1879)
- bullata Körber (1861)
 - Lecidea nigroglomerata* Leighton ex Crombie (1871)
 - carpathica Körber (1865)
 - elaeochroma (Ach.) M. Choisy (1950)
 - elaeochroma f. soralifera (Erichsen) D. Hawksw. (1972)
 - meiococca (Nyl.) Leuckert & Hertel (1990)
 - Lecidea meiococca* Nyl. (1876)
 - Lecidella prasinula* auct., non (Wedd.) Hertel (1980)
- * pulveracea (Schaerer) H. Sydow (1887)
- scabra (Taylor) Hertel & Leuckert (1969)
 - Lecidella prasinula* (Wedd.) Hertel (1980), non auct.
 - stigmataea (Ach.) Hertel & Leuckert (1969)
 - viridans (Flotow) Körber (1855)
 - wulfenii (Hepp) Körber (1861)

Lecidoma G. Schneider & Hertel (1981)

- demissum (Rutström) G. Schneider & Hertel (1981)

Lemmopsis (Vainio) Zahlbr. (1906)

- arnoldiana (Hepp) Zahlbr. (1906)
- ? oblongans (Nyl. ex Crombie) A.L. Sm. (1918)

L. leptogliella (Nyl.) A.L. Sm. (1918) = *Porocyphus*.

Lempholemma Körber (1855)

- botryosum (Massal.) Zahlbr. (1924)
- chalazanum (Ach.) B. de Lesd. (1910)
- cladodes (Tuck.) Zahlbr. (1924)
 - L. albonigrum* Magnusson (1946)
 - intricatum (Arnold) Zahlbr. (1924)
 - L. fennicum* (Räsänen) Degel. (1946)
 - polyanthes (Bernh.) Malme (1918)
 - ? *L. chalazanellum* (Nyl.) Zahlbr. (1924)
 - ? *L. chalazanodes* (Nyl.) Zahlbr. (1924)
 - L. myriococcum* (Ach.) Th. Fr. (1861)
 - radiatum (Sommerf.) Henssen (1968)

Lepraria Ach. (1803)

- caesioalba (B. de Lesd.) Laundon (1992)
- L. zonata* Brodo (1968)
- crassissima (Hue) Lettau (1958)

- eburnea Laundon (1992)
 frigida Laundon (1992)
 incana (L.) Ach. (1803)
 jackii Tønsb. (1992)
lesdainii (Hue) R. Harris (1987)
lobificans Nyl. (1873)
L. finkii (B. de Lesd. ex Hue) R. Harris (1985)
neglecta (Nyl.) Lettau (1958)
? *Lecidea pedatula* Nyl. (1876)
nivalis Laundon (1992)
L. crassissima auct. p.p., non (Hue) Lettau (1958)
? *rigidula* (Hue) Tønsb. (1992)
umbricola Tønsb. (1992)

L. membranacea (Dickson) Vainio (1921) = *Leproloma*

- Leprocaulon** Nyl. ex Lamy (1878)
microscopicum (Vill.) Gams ex D. Hawksw. (1974)

- Leproloma** Nyl. ex Crombie (1894)
cacuminum (Massal.) Laundon (1992)
L. angardianum (Øvstedral) Laundon (1989)
diffusum Laundon (1989)
diffusum var. *chrysodetoides* Laundon (1989)
membranaceum (Dickson) Vainio (1899)
Leparia membranacea (Dickson) Vainio (1921)
vouauxii (Hue) Laundon (1989)

- Leproplaca** (Nyl.) Harm. (1900)
chrysodeta (Vainio ex Räsänen) Laundon (1974)
Caloplaca chrysodera (Vainio ex Räsänen) Dombr. (1970)
xantholyta (Nyl.) Harm. (1900)
Caloplaca xantholyta (Nyl.) Jatta (1902)

- Leptogium** (Ach.) Gray (1821)
biatorinum (Nyl.) Leighton (1879)
L. byssinum auct. brit., non (Hoffm.) Zwackh ex Nyl. (1856)
L. cretaceum (Sm.) Nyl. (1855)
brebissonii Mont. (1840)
britannicum P.M. Jørg. & P. James (1983).
burgessii (L.) Mont. (1840)
byssinum (Hoffm.) Zwackh ex Nyl. (1856)
cochleatum (Dickson) P.M. Jørg. & P. James (1983)
L. azureum auct. brit., non (Swartz) Mont. (1840)
corniculatum (Hoffm.) Minks (1873)
coralloideum (Meyen & Flotow) Vainio (1921).
L. palmatum (Huds.) Mont. (1846)
cyanescens (Rabenh.) Körber (1855)

- diffractum Krempelh. ex Körber (1865)
 gelatinosum (With.) Laundon (1984)
L. sinuatum (Huds.) Massal. (1853)
hibernicum Mitchell ex P.M Jørg. (1973)
 * *hildenbrandii* (Garov.) Nyl. (1856)
imbricatum P.M. Jørg. in ed. (1993)
intermedium (Arnold) Arnold (1885)
juressianum C. Tav. (1950)
lichenoides (L.) Zahlbr. (1924)
massiliense Nyl. (1879)
plicatile (Ach.) Leighton (1879)
saturninum (Dickson) Nyl. (1856)
schraderi (Ach.) Nyl. (1856)
subtile (Schrader) Torss. (1843)
tenuissimum (Dickson) Körber (1855)
L. humosum Nyl. (1858)
teretiusculum (Wallr.) Arnold (1892)
turgidum (Ach.) Crombie (1870)

Literature: Jørgensen (1993)

- Leptorhaphis** Körber (1855) [F]
atomaria (Ach.) Szat. (1928) [F]
epidermidis (Ach.) Th. Fr. (1860) [F]
maggiana (Massal.) Körber (1865) [F]

L. ischnobela (Nyl.) Coppins (1980) = *Celothelium*.

- Leucocarpia** Vězda (1969)
biatorella (Arnold) Vězda (1969)

- Leucocarpopsis** G. Salisb. (1975)
devensis G. Salisb. (1975)

- Lichenothelia** D. Hawksw. (1981) [F]
convexa Henssen (1987) [F]

- Lichina** Agardh (1817)
confinis (Müller) Agardh (1821)
pygmaea (Lightf.) Agardh (1821)

- Lichenodium** Nyl. (1875)
sirosiphoideum Nyl. (1875)

- Lithographa** Nyl. (1857)
tesserata (DC.) Nyl. (1857)

Lobaria (Schreber) Hoffm. (1796)
amplissima (Scop.) Forss. (1883)
Dendriscocaulon umhausense (Auersw.) Degel. (1942) [blue-green
 photosymbiodeme]
pulmonaria (L.) Hoffm. (1796)
scrobiculata (Scop.) DC. (1805)
virens (With.) Laundon (1984)
L. laetevirens Zahlbr. (1906)

Lopadium Körber (1855)
coralloideum (Nyl.) Lynge (1940)
disciforme (Flot.) Kullh. (1870)
pezizoideum (Ach.) Körber (1855)

Loxospora Massal. (1852)
elatina (Ach.) Massal. (1852)
Haematomma elatinum (Ach.) Massal. (1854)

Macentina Vězda (1973)
abscondita Coppins & Vězda (1977)
stigonemoides A. Orange (1989)

M. aurantiaca McCarthy & Vězda (1985) = *Nectria pseudopeziza*.

Massalongia Körber (1855)
carnosa (Dickson) Körber (1855)

Megalaria Hafellner (1984).
grossa (Pers. ex Nyl.) Hafellner (1984)
Catinaria grossa (Pers. ex Nyl.) Vainio (1922).

Megalospora Meyen (1843)
tuberculosa (Fée) Sipman (1983)
Bombyliospora pachycarpa auct., non (Delise ex Duby) Massal. (1852)

Megaspora (Clauz. & Roux) Hafellner & V. Wirth (1987)
verrucosa (Ach.) Hafellner & Wirth (1987).
Pachyospora verrucosa (Ach.) Massal. (1852)

Melanelia Esslinger (1978) - see *Parmelia*.

Melanolecia Hertel in Poelt & Vězda (1981) - see *Farnoldia*.

Melaspilea Nyl. (1856) [\pm L]
 * *amota* Nyl. (1867)
atroides Coppins (1992)
 * *diplasiospora* (Nyl.) Müll. Arg. (1887)
granitophila (Th. Fr.) Coppins (1989)

Arthonia granitophila Th. Fr. (1865)

M. subarenacea J. Nowak & Kiszka (1974)

interjecta (Leighton) A.L. Sm. (1911)

Encephalographa interjecta (Leighton) Th. Fr. (1874)

* *leciographoides* Vouaux (1913) [LF]

lentiginosa (Lyell ex Leighton) Müll. Arg. (1887) [LF]

lentiginosula (Nyl.) A.L. Sm. (1911) [F]

ochrothalamia Nyl. (1865)

proximella Nyl. ex Norrlin (1873) [F]

Menegazzia Massal. (1854)

terebrata (Hoffm.) Massal. (1854)

Micarea Fr. (1825)

adnata Coppins (1983)

alabastrites (Nyl.) Coppins (1982)

assimilata (Nyl.) Coppins (1983)

Lecidea assimilata Nyl. (1861)

bauschiana (Körber) V. Wirth, & Vězda (1976)

Lecidea rusticella Nyl. (1878)

botryoides (Nyl.) Coppins (1980)

cinerea (Schaerer) Hedl. (1892)

Bacidia carneoalbens (Nyl.) A.L. Sm. (1911)

Hastifera tenuispora D. Hawksw. & Poelt (1986) [anamorph]

contexta Hedl. (1892)

coppinsii Tønsb. (1992)

crassipes (Th. Fr.) Coppins (1983)

Helocarpon crassipes Th. Fr. (1860)

denigrata (Fr.) Hedl. (1892)

erratica (Körber) Hertel, Rambold & Pietschm. (1989)

Lecidea erratica Körber (1861)

Lecidea tephrizans Leighton (1876)

excipulata Coppins (1988)

globulosella (Nyl.) Coppins (1983)

M. bacidiella sensu Vězda, non (Vainio) Vězda (1976)

granulans (Vainio) Timdal (1991)

M. subleprosula (Vězda) Vězda (1976)

hedlundii Coppins (1983)

incrassata Hedl. (1892)

inquinans (Tul.) Coppins (1992)

Lecidea inquinans (Tul.) Nyl. (1856)

intrusa (Th. Fr.) Coppins & Kilias (1983)

Catillaria intrusa (Th. Fr.) Th. Fr. (1874)

Lecidea aphanoides Nyl. (1868)

Lecidea melaphana Nyl. (1869)

leprosula (Th. Fr.) Coppins & A. Fletcher (1975)

lignaria (Ach.) Hedl. (1892)

lignaria var. *endoleuca* (Leighton) Coppins (1983)

- lignaria* f. *gomphillacea* (Nyl.) Hedl. (1892)
lithinella (Nyl.) Hedl. (1892)
lutulata (Nyl.) Coppins (1980)
Lecidea botryiza Nyl. ex Stirton (1873)
melaena (Nyl.) Hedl. (1892)
misella (Nyl.) Hedl. (1892)
myriocarpa V. Wirth & Vězda ex Coppins (1983)
nigella Coppins (1983)
nitschkeana (Lahm ex Rabenh.) Harm. (1899)
olivacea Coppins (1983)
peiliocarpa (Anzi) Coppins & R. Sant. (1979)
prasina Fr. (1825)
M. subviridescens (Nyl.) Hedl. (1892)
pseudomarginata Coppins (1988)
pycnidiophora Coppins & P. James (1979)
stipitata Coppins & P. James (1979)
submoestula (Nyl.) Coppins (1992)
Lecidea submoestula Nyl. (1876)
subnigrata (Nyl.) Coppins & Kiliaš (1980)
Catillaria subnigrata (Nyl.) Herre (1910)
Micarea confusa (Nyl.) Hedl. (1892)
subviolascens (Magnusson) Coppins (1983)
sylvicola (Flotow) Vězda & V. Wirth (1976)
synotheoides (Nyl.) Coppins (1982)
ternaria (Nyl.) Vězda (1970)
tuberculata (Sommerf.) R. Anderson (1974)
turfosa (Massal.) Du Rietz (1923)
viridiatra Coppins (1985)

M. clavulifera (Nyl.) Coppins & P. James (1980) = *Psilolechia*.

- Microcalicum** Vainio (1927) [F/LF]
ahlneri Tibell (1978) [F]
arenarium (Hampe ex Massal.) Tibell (1978) [LF]
disseminatum (Ach.) Vainio (1927) [F]
M. subpedicillatum (Schaerer) Tibell (1978)

- Microglaena** Körber (1855)
M. corrosa (Körber) Arnold (1877) = *Protothelenella*
M. larbalestieri A.L. Sm. (1911) = *Chromatochlamys*
M. leucothelia (Anzi) Arnold (1870) = *Protothelenella*
M. modesta (Nyl.) A.L. Sm. (1911) = *Thielenella*
M. muscorum (Fr.) Th. Fr. (1860) = *Chromatochlamys*
M. sphinctrinoidella (Nyl.) Arnold (1870) = *Protothelenella*
M. sphinctrinoides (Nyl.) Lönnr. (1858) = *Protothelenella*.

Microthelia Körber (1855) - see *Mycomicrothelia*.

Miriquidica Hertel & Rambold (1987)

- atrofulva (Sommerf.) Rambold & Schwab (1990)
 - Lecidea atrofulva* Sommerf. (1826)
- complanata (Körber) Hertel & Rambold (1987)
 - Aspicilia complanata* (Körber) B. Stein (1879)
 - Aspicilia superiuscula* (Nyl.) Hue (1912)
- garovagliae (Schaerer) Hertel & Rambold (1987)
- griseoatra (Flotow) Hertel & Rambold (1987)
 - Lecidea griseoatra* (Flotow) Schaerer (1850)
 - ? *Lecidea subplumbea* Anzi (1868)
- leucophaea (Rabenh.) Hertel & Rambold (1987)
 - Lecidea leucophaea* (Rabenh.) Nyl. (1870)
- lulensis (Hellbom) Hertel & Rambold (1987)
 - Lecidea lulensis* Hellbom (1866)
- nigroleprosa (Vainio) Hertel & Rambold (1987)
 - Lecidea nigroleprosa* (Vainio) Magnusson (1927)

Mniacea Boud. (1885) [F]

- jungermanniae* (Nees ex Fr.) Boud. (1906) [F]
- nivea* (Crouan) Boud. (1907) [F]

Moelleropsis Gyelnik (1940)

- humida (Kullh.) Coppins & P.M. Jørg. (1993)
 - Biatora humida* Kullh. (1871)
- nebulosa (Hoffm.) Gyelnik (1904)

Mosigia Fr. (1845) - see *Rimularia*.

Multiclavula R.H. Petersen (1967)

- vernalis* (Schw.) R.H. Petersen (1967)

Mycobilimbia Rehm (1890) - see *Biatora*.

Mycoblastus Norman (1852)

- affinis* (Schaerer) Schauer (1964)
- alpinus* (Fr.) Kernst. (1893)
- caesius* (Coppins & P. James) Tønsb. (1992)
 - Haematomma caesium* Coppins & P. James (1978)
 - fucatus* (Stirton) Zahlbr. (1926)
 - sanguinarius* (L.) Norman (1853)
 - sanguinarius* f. *leprerosus* Nádv. (1951)
 - sterilis* Coppins & P. James (1979)

M. tornoensis (Nyl.) R. Anderson (1974) = *Japewia*.

Mycocalicium Vainio (1890) [F]

- subtile* (Pers.) Szat. (1925) [F]

Mycoglaena Höhnel (1909) [F]

acuminans (Nyl.) Vainio (1921) [F]
myricae (Nyl.) R. Harris (1973) [F]

Mycomicrothelia Keissler (1936) [\pm L]

atlantica D. Hawksw. & Coppins (1985) [F]
confusa D. Hawksw. (1985)

Microthelia micula auct. brit., non Körber (1855)
wallrothii (Hepp) D. Hawksw. (1980) [F]

Mycoporum Flotow ex Nyl. (1855) [F]

hippocastani (DC.) Coppins (1979) [F]
quercus (Massal.) Müll. Arg. (1882) [F]

Neofuscelia Esslinger (1978) - see *Parmelia*.

Nephroma Ach. (1810)

arcticum (L.) Torss. (1843)
* helveticum Ach. (1810)
laevigatum Ach. (1814)
parile (Ach.) Ach. (1810)
* resupinatum (L.) Ach. (1810)
tangeriense (Maheu and A. Gillet) Zahlbr. (1932)

Normandina Nyl. (1855)

pulchella (Borrer) Nyl. (1861)

Ochrolechia Massal. (1852)

androgyna (Hoffm.) Arnold (1885)
. frigida (Swartz) Lynge (1928)
? *O. lapuensis* (Vainio in Räsänen) Räsänen (1931)
? *O. frigida* f. *lapuensis* (Vainio) Coppins (1981)
inaequatula (Ny.) Zahlbr. (1913)
inversa (Nyl.) Laundon (1963)
microstictoides Räsänen (1936)
parella (L.) Massal. (1852)
subviridis (Hoeg) Erichsen (1930)
szatalaënsis Vers. (1958)
O. pallescens auct. brit. p.p., non (L.) Massal. (1853)
tartarea (L.) Massal. (1852)
turneri (Sm.) Hasselrot (1945)

O. geminipara (Th. Fr.) Vainio (1881) = *Pertusaria*.

Omphalina Quélet (1886) [\pm L]

- cupulatoides P.D. Orton (1977) [LF]
 ericetorum (Fr.) M. Lange ex H. Bigelow (1970)
Botrydina vulgaris Bréb. (1839)
 ? *O. fulvopallens* P.D. Orton (1984)
griseopallida (Desm.) Quélet (1886)
hudsoniana (Jennings) H. Bigelow (1970)
Coriscium viride (Ach.) Vainio (1890)
O. luteolilacina (Favre) Henderson (1958)
luteovitellina (Pilát & Nannf.) M. Lange (1957)
pararustica Cléménçon (1982)
pseudoandrosacea (Bull.) Moser (1967)
velutina (Quélet) Quélet (1886)
O. grisella (Weinm.) Moser (1953)

Opegrapha Humb. (1793) [\pm L]

- areniseda Nyl. (1875)
O. actophila Nyl. (1880)
 atra Pers. (1794)
 brevis Coppins (1987) [LF]
cesareensis Nyl. (1868)
O. hysteriformis Nyl. (1876)
corticola Coppins & P. James (1979)
demutata Nyl. (1879)
dolomitica (Arnold) Clauzade & Roux (1989)
O. saxicola auct., non Ach. (1814)
fumosa Coppins & P. James (1992)
gyrocarpa Flotow (1825)
herbarum Mont. (1833)
 ? *O. protuberans* Zahlbr. (1932)
lithyrga Ach. (1810)
 ? *monspeliensis* Nyl. (1857) [LF]
mougeotii Massal. (1853)
multipuncta Coppins & P. James (1992)
niveoatra (Borrer) Laundon (1963)
O. dubia Leighton ex Arnold (1861)
ochrocheila Nyl. (1865)
parasitica (Massal.) H. Olivier (1906) [LF]
O. persoonii (Ach.) Chevall. (1822), nom. illegit.
 ? = *O. rupestris* Pers. (1794)
paraxanthodes Nyl. (1879)
pertusariicola Coppins & P. James (1979) [LF]
physciaria (Nyl.) D. Hawksw. & Coppins (1992) [LF]
prosodea Ach. (1803)
pulvinata Rehm (1869) [LF]
 ? *tubefacta* Räsänen (1946)
rufescens Pers. (1794)

saxatilis auct., non DC. (1805)
 ? *O. calcarea* Turner ex Sm. (1807)
 ? *O. chevallieri* Leighton (1854)
 ? *O. conferta* Anzi (1862)
 sasicola auct., non Ach. (1814)
 saxigena Tayl. (1836)
sorediifera P. James (1962)
subelevata Nyl. (1888)
thelotrematis Coppins (1984) [LF]
varia Pers. (1794)
O. atrula Nyl. (1877)
O. diaphora (Ach.) Ach. (1803)
O. lichenoides Pers. (1794)
O. rimalis Ach. (1809)
vermicellifera (Kunze) Laundon (1963)
viridis (Ach.) Nyl. (1861)
vulgata (Ach.) Ach. (1803)
O. cineréa Chevall. (1822)
xerica Torrente & Egea (1992)

O. lyncea (Sm.) Borrer ex Hook. (1833) = *Lecanactis*
O. zonata Körber (1855) = *Enterographa*.

Ophioparma Norman (1853)
ventosum (L.) Norman (1853)
Haematomma ventosum (L.) Massal. (1852)

Orphniospora Körber (1874)
moriopsis (Massal.) D. Hawksw. (1982)
O. atrata (Sm.) Poelt (1972)
moriopsis var. *brunnea* (A.L. Sm) D. Hawksw. (1982)
O. atrata var. *brunnea* (A.L. Sm.) Poelt (1980)

Pachyphiale Lönnr. (1858)
carneola (Ach.) Arnold (1871)
P. cornea (Ach.) Poetsch (1872)
fagicola (Hepp) Zwackh (1862)

Pachyospora auct., non Massal. (1852) - see *Megaspóra*.

Pannaria Delise (1828)
conoplea (Ach.) Bory (1828)
hookeri (Borrer ex Sm.) Nyl. (1857)
ignobilis Anzi (1862)
leucophaea (Vahl) P.M. Jørg. (1978)
mediterranea C. Tav. (1965)
pezizoides (Weber) Trevisan (1869)

- praetermissa* Nyl. (1858)
Parmeliella praetermissa (Nyl.) P. James (1965)
ruginosa (Ach.) Bory (1828)
sampaiana C. Tav. (1950)
- Parmelia** Ach. (1803)
acetabulum (Necker) Duby (1830)
Pleurosticta acetabulum (Necker) Elix & Lumbsch (1988)
arnoldii Du Rietz (1924)
Parmotrema arnoldii (Du Rietz) Hale (1974)
borreri (Sm.) Turner (1807)
Punctelia borreri (Sm.) Krog (1982)
britannica D. Hawksw. & P. James (1973)
caperata (L.) Ach. (1803)
Flavoparmelia caperata (L.) Hale (1986)
conspersa (Ehrh. ex Ach.) Ach. (1803)
Xanthoparmelia conspersa (Ehrh. ex Ach.) Hale (1974)
crinita Ach. (1814)
Parmotrema crinitum (Ach.) M. Choisy (1952)
delisei (Duby) Nyl. (1872)
Neofuscelia delisei (Duby) Essl. (1978)
discordans Nyl. (1886)
P. omphalodes subsp *discordans* (Nyl.) Skult (1984)
disjuncta Erichsen (1939)
Melanelia disjuncta (Erichsen) Essl. (1978)
elegantula (Zahlbr.) Szat. (1930)
Melanelia elegantula (Zahlbr.) Essl. (1978)
endochlora Leighton (1871)
Hypotrichyna endochlora (Leighton) Hale (1975)
exasperata de Not. (1847)
Melanelia exasperata Essl. (1978)
exasperatula Nyl. (1873)
Melanelia exasperatula (Nyl.) Essl. (1978)
glabratula (Lamy) Nyl. (1883)
Melanelia glabratula (Lamy) Essl. (1978)
glabratula subsp. *fuliginosa* (Fr. ex Duby) Laundon (1965)
horrescens Taylor (1836)
Parmelinopsis horrescens (Taylor) Elix & Hale (1987)
incurva (Pers.) Fr. (1826)
Arctoparmelia incurva (Pers.) Hale (1986)
laciniatula (H. Olivier) Zahlbr. (1916)
Melanelia laciniatula (H. Olivier) Essl. (1978)
laevigata (Sm.) Ach. (1814)
Hypotrichyna laevigata (Sm.) Hale (1975)
loxodes Nyl. (1872)
Neofuscelia loxodes (Nyl.) Essl. (1978)
minarum Vainio (1890)
Parmelinopsis minarum (Vainio) Elix & Hale (1987)

- mougeotii Schaeer ex D. Dietr. (1846)
Xanthoparmelia mougeotii (Schaeer ex D. Dietr.) Hale (1974)
- omphalodes (L.) Ach. (1803)
pastillifera (Harm.) R. Schubert & Klem. (1966)
Parmelina pastillifera (Harm.) Hale (1976)
- perlata (Huds.) Ach. (1803)
P. coniocarpa Lauter (1827)
Parmotrema chinense (Osbeck) Hale and Ahti (1986)
- protomatrae Gyelnik (1931)
Xanthoparmelia protomatrae (Gyelnik) Hale (1974)
- pulla Ach. (1814)
Neofuscelia pulla (Ach.) Essl. (1978)
quercina (Willd.) Vainio (1899)
Parmelina quercina (Willd.) Hale (1974)
- reddenda Stirton (1877)
Punctelia reddenda (Stirton) Krog (1982)
- reticulata Taylor (1836)
Parmotrema reticulatum (Taylor) M. Choisy (1952)
Rimelia reticulata (Taylor) Hale & A. Fletcher (1990)
- revoluta Flörke (1827)
Hypotrachyna revoluta (Flörke) Hale (1975)
- robusta Degel. (1941)
Parmotrema robustum (Degel.) Hale (1974)
- saxatilis (L.) Ach. (1801)
septentrionalis (Lynge) Ahti (1966)
Melanelia septentrionalis (Lynge) Essl. (1978)
- sinuosa (Sm.) Ach. (1814)
Hypotrachyna sinuosa (Sm.) Hale (1975)
- soredians Nyl. (1872)
Flavoparmelia soredians (Nyl.) Hale (1986)
- stygia (L.) Ach. (1803)
Melanelia stygia (L.) Essl. (1978)
- subargentifera Nyl. (1875)
Melanelia subargentifera (Nyl.) Essl. (1978)
- subaurifera Nyl. (1873)
Melanelia subaurifera (Nyl.) Essl. (1978)
- submontana Nadv. ex Hale (1987)
- subrudecta Nyl. (1888)
Punctelia subrudecta (Nyl.) Krog (1982)
- sulcata Taylor (1836)
taylorensis M. Mitchell (1961)
Hypotrachyna taylorensis (M. Mitchell) Hale (1974)
- tiliacea (Hoffm.) Ach. (1803)
Parmelina tiliacea (Hoffm.) Hale (1974)
- tinctina Maheu & A. Gillet (1925)
Xanthoparmelia tinctina (Maheu & A. Gillet) Hale (1974)
- verruculifera Nyl. (1878)
Neofuscelia verruculifera (Nyl.) Essl. (1978)

Parmeliella Müll. Arg. (1862)

jamesii S. Ahlner & P.M. Jørg. (1978)
 testacea P.M. Jørg. (1978)
 triptophylla (Ach.) Müll. Arg. (1862)

P. atlantica Degel. (1935) = *Degelia*

P. plumbea (Lightf.) Vainio (1890) = *Degelia*

P. praetermissa (Nyl.) P. James (1965) = *Pannaria*

Parmelina Hale (1974) - see *Parmelia*.

Parmelinopsis Elix & Hale (1987) - see *Parmelia*.

Parmeliopsis (Nyl.) Nyl. (1866)

ambigua (Wulfen) Nyl. (1866)

Foraminella ambigua (Wulfen) Fricke Meyer (1982)

hyperopta (Ach.) Arnold (1884)

Foraminella hyperopta (Ach.) Fricke Meyer (1982)

P. aleurites (Ach.) Nyl. (1866) = *Imshaugia*.

Parmotrema Massal. (1860) - see *Parmelia*.

Parmentaria Fée (1825)

chilensis Fée (1838)

? *Pyrenula chilensis* (Fée) R. Harris (1989)

Peltigera Willd. (1787)

britannica (Gyelnik) Holt.-Hartw. & Tønsb. (1983)

canina (L.) Willd. (1787)

collina (Ach.) Schrader (1801)

degenii Gyelnik (1927)

didactyla (With.) Laundon (1984)

P. spuria (Ach.) DC. (1805)

elisabethae Gyelnik (1927)

horizontalis (Huds.) Baumg. (1790)

lactucifolia (With.) Laundon (1984)

P. hymenina (Ach.) Delise ex Duby (1830)

lepidophora (Nyl.) Bitter (1904)

leucophlebia (Nyl.) Gyelnik (1926)

malacea (Ach.) Funck (1827)

membranacea (Ach.) Nyl. (1887)

neckeri Hepp ex Müll. Arg. (1862)

polydactyla (Necker) Hoffm. (1790)

ponojensis Gyelnik (1931)

praetextata (Flörke ex Sommerf.) Zopf (1909)

rufescens (Weis) Humb. (1793)

scabrosa Th. Fr. (1861)

venosa (L.) Hoffm. (1789)

Pertusaria DC. (1805)

- albescens (Huds.) M. Choisy & Werner (1932)
- albescens var. corallina (Zahlbr.) Laundon (1963)
- amara (Ach.) Nyl. (1873)
- amara f. pulvinata (Erichsen) Almb. (1948)
- P. pulvinata* Erichsen (1936)
- amarescens Nyl. (1874)
- aspergilla (Ach.) Laundon (1992)
- P. dealbescens* auct., non Erichsen (1936)
- borealis Erichsen (1938)
- bryontha (Ach.) Nyl. (1861)
- chiodectonoides Bagl. ex Massal. (1856)
- coccodes (Ach.) Nyl. (1857)
- corallina (L.) Arnold (1861)
- P. areolata* auct. brit., non (Ach.) Massal. (1852)
- coronata (Ach.) Th. Fr. (1871)
- dactylna (Ach.) Nyl. (1863)
- excludens Nyl. (1885)
- flavicans Lamy (1878)
- flavida (DC.) Laundon (1963)
- flavocorallina Coppins & Muhr (1992)
- gallica B. de Lesd. (1925)
- geminipara (Th. Fr.) Knight (1883)
- Ochrolechia geminipara* (Th. Fr.) Vainio (1881)
- glomerata (Ach.) Schaeerer (1826)
- hemisphaerica (Flörke) Erichsen (1932)
- hymenea (Ach.) Schaeerer (1836).
- P. rupicola* auct. brit., non (Fr.) Harm. (1898)
- lactea (L.) Arnold (1872)
- * lactescens Mudd (1861)
- leioplaca DC. (1815)
 - ? *P. alpina* Hepp ex Ahles. (1860)
 - P. leucostoma* Massal. (1852)
 - melanochlora (DC.) Nyl. (1872)
 - monogona Nyl. (1872)
 - multipuncta (Turner) Nyl. (1861)
 - oculata (Dickson) Th. Fr. (1871)
 - ophthalmiza (Nyl.) Nyl. (1865)
- * panyrga auct. brit., non (Ach.) Massal. (1855)
 - P. hutchinsiae* (Borrer) Leighton (1851)
 - pertusa (Weigel) Tuck. (1845)
 - ? *P. colliculosa* Körber (1863)
 - ? polythecia (Taylor) Erichsen (1936)
 - pseudocorallina (Lilj.) Arnold (1887)
 - P. ceuthocarpoides* Zahlbr. (1927)

pupillaris (Nyl.) Th. Fr. (1871)

Lecidea cinnabarina auct. brit. p.p., non Sommerf. (1823)

pustulata (Ach.) Duby (1830)

velata (Turner) Nyl. (1861)

xanthostoma (Sommerf.) Fr. (1831)

Aspicilia poriniformis (Nyl.) Arnold (1870)

Petractis Fr. (1846)

clausa (Hoffm.) Krempelh. (1861)

Phaeocalicium A. Schmidt (1970) [F]

populneum (Brond ex Duby) A. Schmidt (1970) [F]

praecedens (Nyl.) A. Schmidt (1970) [F]

Phaeographis Müll. Arg. (1882)

dendritica (Ach.) Müll. Arg. (1882)

inusta (Ach.) Müll. Arg. (1881)

lyelli (Sm.) Zahlbr. (1905)

smithii (Leighton) B. de Lesd. (1910)

Phaeophyscia Moberg (1977)

endococcina Körber) Moberg (1977)

endophoenicea (Harm.) Moberg (1977)

nigricans (Flörke) Moberg (1977)

orbicularis (Necker) Moberg (1977)

Physcia labrata auct. brit., non Meresch (1919)

P. orbicularis var. *hueana* (Harm.) Clauz. & Roux (1985)

sciastra (Ach.) Moberg (1977)

Phaeopyxis Rambold & Triebel (1990) [LF]

punctum (Massal.) Rambold, Triebel & Coppins (1990) [LF]

? *Lecidea cladoniaria* Nyl. (1857)

Lecidea punctum (Massal.) Jatta (1900)

varia Coppins, Rambold & Triebel (1990) [LF]

Phlyctis Wallr. (1825)

agelaea (Ach.) Flotow (1850)

argena (Sprengel) Flotow (1850)

Phyllopsora Müll. Arg. (1894)

rosei Coppins & P. James (1979)

Physcia (Schreber) Michaux (1803)

adscendens (Fr.) H. Olivier (1882)

aipolia (Ehrh. ex Humb.) Fürnrohr (1839)

caesia (Hoffm.) Fürnrohr (1839)

P. subalbinea Nyl. (1874)

P. wainioi Räsänen (1921)

clementei (Sm.) Maas Geest. (1952)

- dubia* (Hoffm.) Lettau (1912)
P. teretiuscula (Ach.) Lyngé (1916)
semipinnata (Gmelin) Moberg (1977)
stellaris (L.) Nyl. (1856)
tenella (Scop.) DC. (1805)
tenella subsp. *marina* (A.E. Nyl.) D. Hawksw. (1980)
tribacia (Ach.) Nyl. (1874)
tribacioides Nyl. (1874)

Physciopsis M. Choisy (1950) - see *Hyperphyscia*.

Physconia Poelt (1965)

- distorta* (With.) Laundon (1984)
Physconia pulverulacea Moberg (1979)
enteroxantha (Nyl.) Poelt (1966)
grisea (Lam.) Poelt (1965)
perisidiosa (Erichsen) Moberg (1977)

Phytoconis Bory (1797) - see *Omphalina*.

Pilophorus Th. Fr. (1857)

- strumaticus* Nyl. ex Crombie (1875)

Placiopsis Beltram. (1858)

- cartilaginea* (Nyl.) Vainio (1921)
P. custnani (Massal.) Körber (1863)
pseudocinerea Breuss (1983)

Placopsis (Nyl.) Lindsay (1866)

- gelida* (L.) Lindsay (1866)
lambii Hertel & V. Wirth (1987)

Placynthiella Elenkin (1909)

- dasaea* (Stirton) Tönsb. (1992)
Lecidea dasaea Stirton (1880)
hyporhoda (Th. Fr.) Coppins & P. James (1984)
? *Lecidea uliginascens* Stirton (1878)
icmalea (Ach.) Coppins & P. James (1984)
oligotropha (Laundon) Coppins & P. James (1984)
Lecidea oligotropha Laundon (1960)
uliginosa (Schrader) Coppins & P. James (1984)
Lecidea uliginosa (Schrader) Ach. (1803)

Placynthium (Ach.) Gray (1821)

- asperellum* (Ach.) Trevisan (1869)
flabellosum (Tuck.) Zahlbr. (1925)
garovaglii (Massal.) Malme (1918)
lismoreense (Nyl. ex Crombie) Vainio (1909)

nigrum (Huds.) Gray (1821)
 pannariellum (Nyl.) Magnusson (1936)
 pluriseptatum (Arnold) Arnold (1889)
 subradiatum (Nyl.) Arnold (1884)
 tantaleum (Hepp) Hue (1906)
 tremniacum (Massal.) Jatta (1900)

Platismatia Culb. & C. Culb. (1968)
 glauca (L.) Culb. & C. Culb. (1968)
 norvegica (Lyngé) Culb. & C. Culb. (1968)

Pleurosticta Petrak (1931) - see *Parmelia*.

Poeltinula Hafellner (1984)
 cerebrina (DC.) Hafellner (1984)
 . *Encephalographa cerebrina* (DC.) Massal. (1856)

Polyblastia Massal. (1852)
 agraria Th. Fr. (1864)
 albida Arnold (1858)
 cruenta (Körber) P. James & Swinscow (1971)
 cupularis Massal. (1852)
 ? *P. bryophila* auct. brit., non Lönnr. (1858)
 deminuta Arnold (1861)
 dermatodes Massal. (1855)
 efflorescens Coppins (1991)
 gelatinosa (Ach.) Th. Fr. (1862)
 * ? gothica Th. Fr. (1865)
 helvetica Th. Fr. (1865)
 inumbrata (Nyl.) Arnold (1870)
 melaspora (Taylor) Zahlbr. (1921)
 quartzina Lyngé (1928)
 sendtneri Krempelh. (1855)
 terrestris Th. Fr. (1860)
 theleodes (Sommerf.) Th. Fr. (1867)
 verrucosa (Ach.) Lönnr. (1848)
 wheldonii Travis (1947)

P. allobata (Stizenb.) Zsch. (1918) = *Agonimia*
P. discrepans Lahm (1867) = *Merismatium discrepans* (Lahm) Triebel (1989) [syn. *M. lopadii*
 auct. brit. p.p.] [F]
P. nigritella (Nyl.) Arnold (1870) = *Merismatium nigritellum* (Nyl.) Vouaux (1913) [F].

Polyblastiopsis Zahlbr. (1903) - see *Julella*.

Polychidium (Ach.) Gray (1821)
 dendriscum (Nyl.) Henssen (1963)
 muscicola (Swartz) Gray (1821)

Polysporina Vězda (1978)

- cyclocarpa (Anzi) Vězda (1978)
 laponica (Schaerer) Degel. (1982)
P. dubia (Magnusson) Vězda (1978)
simplex (Davies) Vězda (1978)

Porina Müll. Arg. (1883)

- aenea (Wallr.) Zahlbr. (1922)
 ahlesiana (Körber) Zahlbr. (1931)
borreri (Trevisan) D. Hawksw. & P. James (1980)
borreri var. *leptospora* (Nyl.) D. Hawksw. (1992)
Porina olivacea var. *leptospora* (Nyl.) Keissler (1937)
bryosiphila (Körber ex Hepp) Zahlbr. (1903)
chlorotica (Ach.) Müll. Arg. (1884)
chlorotica f. *tenuifera* (Nyl.) Swinscow (1962)
curnowii A.L. Sm. (1911)
ginzbergeri Zahlbr. (1903)
grandis (Körber) Zahlbr. (1922)
guentheri (Flotow) Zahlbr. (1922)
guentheri var. *lucens* (Taylor) Swinscow (1962)
heterospora (Fink) R. Harris (1980)
P. nucula auct. brit., non Ach. (1814)
interjungens (Nyl.) Zahlbr. (1922)
lectissima (Fr.) Zahlbr. (1903)
leptalea (Durieu & Mont.) A.L. Sm. (1911)
linearis (Leighton) Zahlbr. (1922)
mammillosa (Th. Fr.) Vainio (1922)
sudetica (Körber) Lettau (1912)

P. coralloidea P. James (1971), *P. hibernica* P. James & Swinscow (1962) and *P. rosei* Sérusiaux (1990) are included in *Zamenhofia*.

Porocyphus Körber (1855)

- coccodes (Flotow) Körber (1855)
kenmorensis (Holl ex Nyl.) Henssen (1974)
leptogiella (Nyl.) L.T. Ellis (1981)
Lemmopsis leptogiella (Nyl.) A.L. Sm. (1918)
rehmicus (Massal.) Zahlbr. (1924)

Porpidia Körber (1855)

- cinereoatra* (Ach.) Hertel & Knoph (1984)
Hulia albocaerulescens auct. brit. p.p., non (Wulfen) Hertel (1975)
Hulia cinereoatra (Ach.) Hertel (1980)
contraponenda (Arnold) Knoph & Hertel (1984)
Hulia albocaerulescens auct. brit. p.p., non (Wulfen) Hertel (1975)
crustulata (Ach.) Hertel & Knoph (1984)
Hulia crustulata (Ach.) Hertel (1975)
Lecidea scutellata W. Watson (1932)

- flavicunda* (Ach.) Gowan (1988)
Huilia flavocaerulescens auct., p.p.
flavocaerulescens (Hornem.) Hertel & Schwab (1984)
Huilia flavocaerulescens (Hornem.) Hertel (1975)
Huilia melinodes (Körber) Hertel (1975)
glaucophaea (Körber) Hertel & Knoph (1984)
Huilia glaucophaea (Körber) Hertel (1875)
hydrophila (Fr.) Hertel & Schwab (1984)
Huilia hydrophila (Fr.) Hertel (1975)
macrocarpa (DC.) Hertel & Schwab (1984)
Huilia macrocarpa (DC.) Hertel (1975)
Huilia nigrocruenta (Anzi) Hertel (1975)
musiva (Körber) Hertel & Knoph (1984)
platycarpoides (Bagl.) Hertel (1987)
Huilia platycarpoides (Bagl.) Hertel (1985)
soredizodes (Lamy ex Nyl.) Laundon (1989)
Huilia soredizodes (Lamy ex Nyl.) Hertel (1980)
speirea (Ach.) Krempelh. (1861)
Lecidea speirea (Ach.) Ach. (1803)
? *L. subumbonella* Lamy (1883)
superba (Körber) Hertel & Knoph (1984)
tuberculosa (Sm.) Hertel & Knoph (1984)
Huilia tuberculosa (Sm.) P. James (1980)

P. athroocarpa (Ach.) Hertel & Rambold (1985) = *Immersaria*.

Protobagliettoa Servít (1955) - see *Verrucaria*.

- Protoblastenia** (Zahlbr.) Steiner (1911)
calva (Dickson) Zahlbr. (1930)
cyclospora (Hepp ex Körber) Poelt (1975)
incrustans (DC.) Steiner (1911)
rupestris (Scop.) Steiner (1911)
siebenhaariana (Körber) Steiner (1911)

- Protoparmelia** M. Choisy (1929)
atriseda (Fr.) R. Sant. & V. Wirth (1986)
Lecanora atriseda (Fr.) Nyl. (1857)
badia (Hoffm.) Hafellner (1984)
Lecanora picea (Dickson) Nyl. (1986)
Lecanora badia (Hoffm.) Ach. (1810)
montagnei (Fr.) Poelt & Nimis (1987)
Lecanora nitens auct., non Ach. (1814)
Lecanora psarophana Nyl. (1872)
nephaea (Sommerf.) R. Sant. (1990)
ochrococca (Nyl.) P.M. Jørg., Rambold & Hertel (1988)
Lecanora ochrococca (Nyl.) Clauz. & Roux (1985)
Lecidea ochrococca Nyl. (1860)

oleagina (Harm.) Coppins (1992)

Lecanora oleagina Harm. (1913)

picea auct.

Lecanora picea auct. p.p., non (Dickson) Nyl. (1868)

Protothelenella Räsänen (1943)

corrosa (Körber) Mayrhofer & Poelt (1985)

Microglaena corrosa (Körber) Arnöld (1877)

? *leucothelia* (Nyl.) Mayrhofer & Poelt (1985)

Microglaena leucothelia (Nyl.) Arnold (1870)

sphinctrinoidella (Nyl.) Mayrhofer & Poelt (1985)

Microglaena sphinctrinoidella (Nyl.) Arnold (1870)

sphinctrinoides (Nyl.) Mayrhofer & Poelt (1985)

Microglaena sphinctrinoides (Nyl.) Lönnr. (1858)

Pseudephebe M. Choisy (1930)

pubescens (L.) M. Choisy (1930)

P. pubescens f. *subciliata* (Nyl.) D. Hawksw. (1977)

Pseudevernia Zopf (1903)

furfuracea (L.) Zopf (1903)

furfuracea var. *ceratea* (Ach.) D. Hawksw. (1969)

Pseudocyphellaria Vainio (1890)

aurata (Ach.) Vainio (1890)

crocata (L.) Vainio (1898)

intricata (Del.) Vainio (1898)

lacerata Degel. (1941)

norvegica (Gyelnik) P. James (1979)

Pseudolecidea Clauz. & Roux (1984), non N. March (1896)

- see *Clauroxia*.

Psilolechia Massal. (1860)

clavulifera (Nyl.) Coppins (1983)

Micarea clavulifera (Nyl.) Coppins & P. James (1980)

leprosa Coppins & Purvis (1987)

lucida (Ach.) M. Choisy (1949)

Psora Hoffm. (1796)

decipiens (Hedwig) Hoffm. (1794)

globifera (Ach.) Massal. (1852)

lurida (Ach.) DC. (1805)

P. petri (Tuck.) Fink (1935)

rubiformis (Ach.) Hook. (1844)

Psoroma Ach. ex Michaux (1803)

hypnorum (Vahl) Gray (1821)

Psorotichia Massal. (1855)

? diffundens (Nyl.) Arnold (1869)
 ? pyrenopsoides (Nyl.) Forss. (1885)
 schaeereri (Massal.) Arnold (1869)

Pterygiopsis Vainio (1890)

coracodiza (Nyl.) Henssen (1990)
 lacustris P.M. Jørg. & R. Sant. (1990)

Ptychographa Nyl. (1874)

flexella (Ach.) Coppins (1980)
 xylographoides Nyl. (1874)

Punctelia Krog (1982) - see *Parmelia*.

Pycnothelia Dufour (1821)

papillaria Dufour (1821)

Pyrenocollema Reinke (1885)

arenisedum (A.L. Sm.) Coppins (1992)
Arthopyrenia areniseda A.L. Sm. (1911)
 argilosporum (Nyl.) Coppins (1992)
Arthopyrenia argilospora (Nyl.) A.L. Sm. (1918)
 bryosporum (Nyl.) Coppins (1992)
Arthopyrenia bryospora (Nyl.) Arnold (1870)
 caesium (Nyl.) R. Harris (1987)
Arthopyrenia caesia (Nyl.) Zahlbr. (1921)
 elegans R. Sant. (1992)
Arthopyrenia elegans R. Sant. (1985), nom. inval.
 halodytes (Nyl.) R. Harris (1987)
Arthopyrenia halodytes (Nyl.) Arnold (1891)
 monense (Wheldon) Coppins (1992)
Arthopyrenia monensis (Wheldon) Zahlbr. (1922)
 orustense (Erichsen) A. Fletcher (1992)
Arthopyrenia orustensis Erichsen (1930)
 pelvetiae (Sutherl.) D. Hawksw. (1988)
Arthopyrenia pelvetiae (Sutherl.) D. Hawksw. (1980)
 saxicola (Massal.) Coppins (1992)
Arthopyrenia saxicola Massal. (1865)
 strontianense (Swinscow) R. Harris (1987)
Arthopyrenia strontianensis Swinscow (1967)
 subarenisedum (G. Salisb.) Coppins (1992)
Arthopyrenia subareniseda G. Salisb. (1953)
 sublitorale (Leighton) R. Harris (1987)
Arthopyrenia sublitoralis (Leighton) Arnold (1891)

Pyrenopsisidium (Nyl.) Forss. (1855) - see *Pyrenopsis*.

Pyrenopsis (Nyl.) Nyl. (1858)

furturea (Nyl.) Th. Fr. (1866)

Pyrenopsidium furfureum (Nyl.) Forss. (1885)*Pyrenopsidium homoeopsis* (Nyl.) Forss. (1885)

grumulifera Nyl. (1861)

P. multispora Coppins (1988)

impolita (Th. Fr.) Forss. (1885)

phylliscella Nyl. (1875)

subareolata Nyl. (1861)

? *P. fuscatula* Nyl. (1858)? *P. haematopsis* auct. brit., non (Sommerf.) Th. Fr. (1860)*P. rhodosticta* auct., non (Taylor) Müll. Arg. (1888)

P. granatina (Sommerf.) Nyl. (1866) and *P. pulvinata* (Schaerer) Th. Fr. (1866) are included in *Euopsis*.

Pyrenula Massal. (1852)

chlorospila Arnold (1887)

coryli Massal. (1852)

Mycopyrenula coryli (Massal.) Vainio (1921)

dermatodes (Borrer) Schaeerer (1850)

laevigata (Pers.) Arnold (1885)

macrospora (Degel.) Coppins & P. James (1980)

? microtheca R. Harris (1989)

nitida (Weigel) Ach. (1814)

nitidella (Flörke ex Schaeerer) Müll. Arg. (1885)

occidentalis (R. Harris) R. Harris (1987)

P. neglecta auct. brit., non R. Harris (1973)**Pyrrhospora** Körber (1855)

quernea (Dickson) Körber (1855).

Racodium Pers. (1797)

rupestre Pers. (1797)

Ramalina Ach. (1810)

calicaris (L.) Fr. (1824)

canariensis Steiner (1904)

R. baltica auct. brit. p.p., ? non Lettau (1912)

chondrina Steiner (1904)

cuspidata (Ach.) Nyl. (1870)

farinacea (L.) Ach. (1810)

fastigiata (Pers.) Ach. (1810)

fraxinea (L.) Ach. (1810)

laceria (With.) Laundon (1984)

R. duriae (de Not.) Bagl. (1879)

pollinaria (Westr.) Ach. (1810)

polymorpha (Lilj.) Ach. (1810)

portuensis G. Samp (1924)
siliquosa (Hudson) A.L. Sm. (1918)
subfarinacea (Nyl. ex Crombie) Nyl. (1872)

Ramonia Stizenb. (1862)
chrysophaea (Pers.) Vězda (1966)
dictyospora Coppins (1987)
luteola auct brit., non Vězda (1967)
nigra Coppins (1987)

Rhaphidicyrtis Vainio (1921)
trichosporella (Nyl.) Vainio (1921)

Rhizocarpon Lam. ex DC. (1805)
advenulum (Leighton) Hafellner & Poelt (1976) [LF]
alpicola (Anzi) Rabenh. (1861)
? *amphibium* (Fr.) Th. Fr. (1874)
badioatrum (Flörke ex Sprengel) Th. Fr. (1874)
caeruleoalbum (Krempelh.) Zahlbr. (1927)
chioneum (Norman) Th. Fr. (1874)
cinereovirens (Müll. Arg.) Vainio (1922)
concentricum (Davies) Beltr. (1858)
R. perlutum (Nyl.) Zahlbr. (1905)
copelandii (Körber) Th. Fr. (1874)
distinctum Th. Fr. (1874)
eupetraeoides (Nyl.) Blomb. & Forss. (1880)
expallescens Th. Fr. (1874)
? *ferax* Magnusson (1948)
? *R. drepanodes* Feuerer (1978)
furfurosum Magnusson & Poelt (1955)
geminatum Körber (1855)
geographicum (L.) DC. (1805)
? *R. frigidum* Räsänen (1944)
? *R. geographicum* subsp. *arcticum* (Runemark) Hertel (1976)
? *R. geographicum* subsp. *diabasicum* (Räsänen) Poelt & Vězda (1980)
? *R. geographicum* subsp. *frigidum* (Räsänen) Hertel (1976)
? *R. geographicum* subsp. *prospectans* (Räsänen) D. Hawksw. & Sowter (1969)
R. riparium Räsänen (1942)
R. riparium subsp. *lindsayanum* (Räsänen) Thomson (1967)
hochstetteri (Körber) Vainio (1922)
inarense (Vainio) Vainio (1898)
intermediellum Räsänen (1943)
jemtlandicum (Malme) Malme (1914)
lavatum (Fr.) Hazsl. (1884)
lecanorinum Anders (1923)
obscuratum (Ach.) Massal. (1852)
ochrolechia (Poelt & Nimis) Hafellner (1992)

- oederi (Weber) Körber (1861)
 plicatile (Leighton) A.L. Sm. (1911)
R. rubescens Th. Fr. (1874)
 polycarpum (Hepp) Th. Fr. (1874)
 postumum (Nyl.) Arnold (1870)
 richardii (Nyl.) Zahlbr. (1926)
R. constrictum Malme (1919)
 simillimum (Anzi) Lettau (1912)
 subgeminatum Eitner (1911)
 superficiale (Schaerer) Malme (1914)
 umbilicatum (Ramond) Flagey (1894)
 viridiatrum (Wulfen) Körber (1855)

Ropalospora Massal. (1860) - see *Fuscidea*.

Rimelia Hale & Fletcher 1990 - see *Parmelia*.

Rimularia Nyl. (1868)

- badioatra (Krempelh.) Hertel & Rambold (1990)
Lecanora contracta (Th. Fr.) Zahlbr. (1928)
Lecidea illita Nyl. (1879)
Mosigia illita (Nyl.) R. Sant. (1984)
 furvella (Nyl. ex Mudd) Hertel & Rambold (1987)
Lecidea furvella Nyl. ex Mudd (1861)
 fuscosora Muhr & Tønsb. (1989)
 gyrizans (Nyl.) Hertel & Rambold (1990)
Lecidea fuscocinerea auct., non Nyl. (1852)
 insularis (Nyl.) Rambold & Hertel (1985)
Lecidea insularis Nyl. (1852)
 intercedens (Magnusson) Coppins (1993)
Lecanora intercedens Magnusson (1930)
Mosigia intercedens (Magnusson) R. Sant. (1984)
 limborina Nyl. (1868)
Lecidea limborina (Nyl.) Lamy (1878)
Lecidea subgyratula Nyl. (1873)
 mullensis (Stirton) Coppins (1993)
Lecidea mullensis Stirton (1877)
 sphacelata (Th. Fr.) Hertel & Rambold (1990)

Rinodina (Ach.) Gray (1821)

- aspersa (Borrer) Laundon (1986)
Buellia aspersa (Borrer) P. James (1965)
 atrocinerea (Hook.) Körber (1855)
 bilobulata (Nyl.) Sheard (1967)
 bischoffii (Hepp) Massal. (1855)
 calcarea (Arnold) Arnold (1879)
 colobina (Ach.) Th. Fr. (1871)
 confragosa (Ach.) Körber (1855)

conradii Körber (1855)
efflorescens Malme (1927)
exigua Gray (1821)
gennarii Bagl. (1861)
griseosoralifera Coppins (1989)
immersa (Körber) Zahlbr. (1906)
interpolata (Stirton) Sheard (1973)
isidioides (Borrer) Olivier (1909)
lecidinea Mayrh. & Poelt (1979)
luridescens (Anzi) Arnold (1872)
milvina (Wahlenb.) Th. Fr. (1860)
mniarea (Ach.) Körber (1855) var. *cinnamomea* Th. Fr. (1860)
occulta (Körber) Sheard (1967)
oculariopsis Mayrh. (1984)
oxydata (Massal.) Massal. (1854)
parasitica Mayrh. & Poelt (1979)
pyrina (Ach.) Arnold (1881)
roboris (Duf. ex Nyl.) Arnold (1881)
sophodes (Ach.) Massal. (1852)
subglaucescens (Nyl.) Sheard (1967)
teichophila (Nyl.) Arnold (1863)

Roccella DC. (1805)

fuciformis (L.) DC. (1805)
phycopsis Ach. (1810)

Sagiolechia Massal. (1854)

protuberans (Ach.) Massal. (1854)
rhexoblephara (Nyl.) Zahlbr. (1913)

Sarcogyne Flotow (1851)

clavus (DC.) Krempelh. (1861)
privigna (Ach.) Massal. (1854)
? pruinosa auct.
regularis Körber (1855)

Sarcopyrenia Nyl. (1858)

cylindrospora (P. Crouan & H. Crouan) M.B. Aguirre (1990)
gibba (Nyl.) Nyl. (1858)

Sarcosagium Massal. (1856)

campestre (Fr.) Poetsch & Schied. (1872)
campestre var. *macrosporum* Coppins & P. James (1979)

Schadonia Körber (1859)

fecunda (Th. Fr.) Vězda & Poelt (1980)

Schaereria Th. Fr. (1855)

- cineroufa* (Schaerer) Th. Fr. (1861)
corticola Muhr & Tønsb. (1992)
fuscocinerea (Nyl.) Clauzade & Roux (1985)
Aspicilia complanatooides (A.L. Sm.) W. Watson (1935)
Lecidea fuscocinerea Nyl. (1852), non auct. brit.
S. endocyanea (Stirton) Hertel & G. Schneider (1980)
S. tenebrosa (Flotow) Hertel & Poelt (1980)
fuscocinerea var. *sorediata* (Houmeau & Roux) Coppins (1992)

Schismatomma Flotow & Körber ex Massal. (1852)

- cretaceum* (Hue) Laundon (1984)
S. virginium D. Hawksw. & P. James (1974)
decolorans (Turner & Borrer ex Sm.) Clauz. & Vězda (1965)
graphidoides (Leighton) Zahlbr. (1919)
? *S. abietinum* (Humb.) Massal. (1852)
? *S. periculum* (Ach.) Branth & Rostrup (1869)
niveum D. Hawksw. & P. James (1971)
quercicola Coppins & P. James (1989)
umbrinum (Coppins & P. James) P.M. Jørg. & Tønsb. (1988)
Lecanactis umbrina Coppins & P. James (1979)

Sclerophora Chevall. (1826)

- nivea* (Hoffm.) Tibell (1984)
Coniocybe pallida (Pers.) Fr. (1824)
peronella (Ach.) Tibell (1984)
Coniocybe peronella (Ach.) Tibell (1978)

Sclerophyton Eschw. (1824)

- circumscripturn* (Taylor) Zahlbr. (1905)

Scoliosporum Massal. (1852)

- chlorococcum* (Graewe ex Stenhammar) Vězda (1978)
pruinose (P. James) Vězda (1978)
sarothamni (Vainio) Vězda (1978)
umbrinum (Ach.) Arnold. (1871)

Scutula Tul. (1852), non Loureiro (1790) - see *Catillaria*.**Siphula** Fr. (1831)

- ceratites* (Wahlenb.) Fr. (1831)

Solenopsora Massal. (1855)

- candicans* (Dickson) Steiner (1915)
holophaea (Mont.) G. Samp. (1921)
liparina (Nyl.) Zahlbr. (1919)
vulturiensis Massal. (1856)

Solorina Ach. (1808)

- bispora Nyl. (1860)
- bispora var. macrospora (Harm.) H. Olivier (1921)
- bispora var. monospora (Gyelnik) Frey (1952)
- crocea (L.) Ach. (1810)
- saccata (L.) Ach. (1810)
- spongiosa (Ach.) Anzi (1862)

Sphaerophorus Pers. (1794)

- fragilis (L.) Pers. (1794)
- globosus (Huds.) Vainio (1903)
- melanocarpus (Swartz) DC. (1815)

Sphinetrina Fr. (1828) [LF]

- anglica Nyl. (1860) [LF]
- leucopoda Nyl. (1860) [LF]
- tubiformis Massal. (1853) [LF]
- turbinata (Pers.) de Not. (1846) [LF]

Spilonema Bornet (1856)

- paradoxum Bornet (1856)
- revertens Nyl. (1865)

Sporastatia Massal. (1854)

- polyspora (Nyl.) Grumm. (1963)
- testudinea (Ach.) Massal. (1854)

Squamaria Poelt (1958)

- cartilaginea (With.) P. James (1980)
- ? cartilaginea var. pseudocrassa (Mattick) D. Hawksw. (1980)
- lentigera (Weber) Poelt (1958)

Staurothele Norman (1853)

- arctica (Lyngé) (1937)
- areolata (Ach.) Lettau (1912)
- S. clopima* auct., non (Wahlenb.) Th. Fr. (1881)
- bacilligera (Arnold) Arnold (1885)
- caesia (Arnold) Arnold (1885)
- fissa (Taylor) Zwackh (1862)
- S. catalepta* sensu Malme (1919), non (Ach.) Blomb. & Forss. (1880)
- geoica Zsch. (1918)
- guestphalica (Lahm ex Körber) Arnold (1885)
- hymenogonia (Nyl.) Th. Fr. (1865)
- rufa (Massal.) Zsch. (1913)
- rugulosa (Massal.) Arnold (1897)
- rupifraga (Massal.) Arnold (1880)
- succedens (Rehm ex Arnold) Arnold (1880)

Literature: Thomson (1991)

Steinia Körber (1873)

geophana (Nyl.) B. Stein (1879)

Stenocybe Nyl. ex Körber (1852) [F]

bryophila W. Watson (1925) [F]

pullatula (Ach.) B. Stein (1879) [F]

S. byssacea (Fr.) Körber (1855)

septata (Leighton) Massal. (1860) [F]

tremulicola Norrlin ex Nyl. (1883) [F]

Stereocaulon Hoffm. (1796)

alpinum Laurer (1827)

condensatum Hoffm. (1796)

dactylophyllum Flörke (1819)

dactylophyllum var. occidentale (Magnusson) Grumm. (1963)

delisei Bory ex Duby (1830)

evolutum Graewe (1865)

glareosum (Savicz) Magnusson (1928)

leucophaeopsis (Nyl.) P. James & Purvis (1985)

Toninia leucophaeopsis (Nyl.) Th. Fr. (1874)

nanodes Tuck. (1859)

pileatum Ach. (1810)

saxatile Magnusson (1926)

spathuliferum Vainio (1909)

symphycheilum Lamb (1961)

* tomentosum Fr. (1824)

tornense (Magnusson) P. James & Purvis (1985)

vesuvianum Pers. (1811)

vesuvianum var. nodulosum (Wallr.) Lamb. (1969)

vesuvianum var. symphycheileoides Lamb in Vězda (1975)

Sticta (Schreber) Ach. (1803)

canariensis (Bory) Bory ex Delise (1823)

S. dufourii Delise (1823) [blue-green photosymbiodeme]

fuliginosa (Hoffm.) Ach. (1803)

limbata (Sm.) Ach. (1803)

sylvatica (Huds.) Ach. (1803)

Strangospora Körber (1860)

delitescens (Arnold) Coppins (1992)

deplanata (Almq.) Clauz. & Roux (1985)

microhaema (Norman) R. Anderson (1975)

moriformis (Ach.) B. Stein (1879)

ochrophora (Nyl.) R. Anderson (1975)

pinicola (Massal.) Körber (1860)

Strigula Fr. (1823)

- jamesii* (Swinscow) R. Harris (1980)
 * *stigmatella* (Ach.) R. Harris (1980)
stigmatella var. *alpestris* (Vězda) Coppins (1980)
taylorii (Carroll ex Nyl.) R. Harris (1980)

Synalissa Fr. (1825)

- symploca* (Ach.) Nyl. (1856)
S. ramulosa (Hoffm.) Körber (1855)

Teloschistes Norman (1853)

- chrysophthalmus* (L.) Th. Fr. (1860)
flavicans (Swartz) Norman (1853)

Tephromela M. Choisy (1929)

- aglaea* (Sommerf.) Hertel & Rambold (1985)
Lecidea aglaea Sommerf. (1826)
armeniaca (DC.) Hertel & Rambold (1985)
Lecidea armeniaca (DC.) Fr. (1825)
atra (Huds.) Hafellner ex Kalb (1983)
Lecanora atra (Huds.) Ach. (1810)
grumosa (Pers.) Hafellner & Roux (1985)
Lecanora grumosa (Pers.) Du Rietz (1926)
pertusarioides (Degel.) Hafellner & Roux (1985)

Thamnolia Ach. ex Schaerer (1850)

- vermicularis* var. *subuliformis* (Ehrh.) Schaerer (1850)

Thelenella Nyl. (1855)

- modesta* (Nyl.) Nyl. (1855)
Microglaena modesta (Nyl.) A.L. Sm. (1911)

Thelidium Massal. (1855)

- aeneovinosum* Anzi (Arnold) (1868)
decipiens (Nyl.) Krempelh. (1861)
? *fontigenum* Massal. (1856)
fumidum (Nyl.) Hazsl. (1884)
T. auruntii auct. brit., non (Massal.) Krempelh. (1861)
impressum (Stizenb.) Zsch. (1920)
incavatum Mudd (1861)
microbolum (Tuck. in Drummond) Hasse (1913)
minutulum Körber (1863)
T. hospitum Arnold (1882)
T. margaceum (Leighton) Zsch. (1920)
T. mesotropum (Nyl.) A. L. Sm. (1911)
T. terrestre W. Watson (1917)
papulare (Fr.) Arnold (1885)
papulare f. *sorediatum* Coppins (1991)

- pluvium A. Orange (1991)
pyrenophorum (Ach.) Mudd (1861)
T. auruntii (Massal.) Krempelh. (1861)
zwackhii (Hepp) Massal. (1855)
T. microcarpum (Davies ex Leighton) A. L. Sm. (1911)
T. subgelatinosum (Sm.) Zsch. (1920)

Thelocarpon Nyl. (1853) [\pm L]

- epibolum Nyl. (1866)
epibolum var. epithallinum (Leighton) G. Salisbury (1953)
impressellum Nyl. (1867)
intermediellum Nyl. (1865)
laureri (Flotow) Nyl. (1855)
lichenicola (Fuckel) Poelt & Hafellner (1975) [LF]
magnussonii G. Salisb. (1953)
olivaceum B. de Lesd. (1914)
pallidum G. Salisb. (1953)
? strasseri Zahlbr. (1902)
superellum Nyl. (1866)

Thelomma Massal. (1860)

- ocellatum (Körber) Tibell (1976)

Thelopsis Nyl. (1855)

- melathelia Nyl. (1864)
rubella Nyl. (1855)

Thelotrema Ach. (1803)

- * isidioides (Borrer) R. Sant. (1980)
lepadinum (Ach.) Ach. (1803)
monosporum Nyl. (1861)
subtile (Tuck.) (1858)

Thermutis Fr. (1825)

- velutina (Ach.) Flotow (1850)

Thrombium Wallr. (1831)

- epigaeum (Pers.) Wallr. (1831)
thelestomum (Ach. ex Harriman) A.L. Sm. (1911)

Tomasellia Massal. (1856) [F]

- diffusa (Leighton) Lahm (1885) [F]
gelatinosa (Chevall.) Zahlbr. (1922) [F]
lactea (Ach.) R. Harris (1980) [F]

Toninia Massal. (1852) [\pm L]

- aromatica (Sm.) Massal. (1852)
? *T. fusispora* (Körber) Th. Fr. (1867)

- cumulata* (Sommerf.) Th. Fr. (1874)
diffracta (Massal.) Zahlbr. (1901)
episema (Nyl.) Timdal (1991) [LF]
Catillaria episema (Nyl.) H. Olivier (1905)
Kiliasia episema (Nyl.) Hafellner (1984)
Scutula episema (Nyl.) Zopf (1896)
lobulata (Sommerf.) Lynge (1926)
mesoidea (Nyl.) Zahlbr. (1926)
opuntioides (Vill.) Buschardt (1979)
* *oribata* (Nyl.) P. James (1965)
physaroides (Opiz) Zahlbr. (1926)
T. lurida (Arnold) H. Olivier (1911)
plumbina (Anzi) Hafellner & Timdal (1991) [LF]
Bacidia plumbina (Anzi) R. Sant. (1960)
* *rosulata* (Anzi) H. Olivier (1911)
sedifolia (Scop.) Timdal (1991)
T. caeruleonigricans (Lightf.) Th. Fr. (1874)
squalescens (Nyl.) Th. Fr. (1874)
squalida (Ach.) Massal. (1852)
T. havaasii Magnusson (1949)
thiopsora (Nyl.) H. Olivier (1911)
T. pulvinata (Taylor) H. Olivier (1911)
tumidula (Sm.) Zahlbr. (1926)
verrucarioides (Nyl.) Timdal (1991)
T. cervina Lönnr. (1858)
T. kolax Poelt (1957)

T. leucophaeopsis (Nyl.) Th. Fr. (1874) = *Stereocaulon*.

- Tornabea** Østh. (1980)
* *scutellifera* (With.) Laundon (1984)
Tornabenia atlantica (Ach.) Kurok. (1962)

- Trapelia** M. Choisy (1929)
coarctata (Sm.) M. Choisy (1932)
corticola Coppins & P. James (1984)
involuta (Taylor) Hertel (1973)
mooreana (Carroll) P. James (1975)
obtegens (Th. Fr.) Hertel (1970)
placodiooides Coppins & P. James (1984)

- Trapeliopsis** Hertel & G. Schneider (1980)
aeneofusca (Flörke ex Flotow) Coppins & P. James (1984)
flexuosa (Fr.) Coppins & P. James (1984)
Lecidea aeruginosa Borrer (1831)
gelatinosa (Flörke) Coppins & P. James (1984)
Lecidea gelatinosa Flörke (1809)
glaucolepidea (Nyl.) G. Schneider (1980)

granulosa (Hoffm.) Lumbsch (1983)

Lecidea granulosa (Hoffm.) Ach. (1803)

percrenata (Nyl.) G. Schneider (1980)

pseudogranulosa Coppins & P. James (1984)

viridescens (Schrader) Coppins & P. James (1984)

Lecidea viridescens (Schrader) Ach. (1803)

wallrothii (Flörke ex Sprengel) Hertel & G. Schneider (1980)

Tremolecia M. Choisy (1953)

atrata (Ach.) Hertel (1977)

T. jurana (Schaerer) Hertel (1977) = *Farnoldia*.

Trimmatothele Norman ex Zahlbr. (1903)

perquisita (Norman) Norman ex Zahlbr. (1903)

Tuckermannopsis Gyelnik (1933) - see *Cetraria*.

Tylothallia P. James & Kilias (1981)

biformigera (Leighton) P. James & Kilias (1981)

Catillaria biformigera (Leighton) Magnusson (1949)

Umbilicaria Hoffm. (1789)

crustulosa (Ach.) Frey (1931)

cylindrica (L.) Delise ex Duby (1830)

deusta (L.) Baumg. (1790)

grisea Hoffm. (1796)

hirsuta (Swartz ex Westr.) Hoffm. (1796)

hyperborea (Ach.) Hoffm. (1796)

nylanderiana (Zahlbr.) Magnusson (1937)

polyphylla (L.) Baumg. (1790)

polyrrhiza (L.) Fr. (1825)

proboscidea (L.) Schrader (1794)

spodochroa (Ehrh. ex Hoffm.) DC. (1805)

torrefacta (Lightf.) Schrader (1794)

Usnea Hill (1753)

articulata (L.) Hoffm. (1796)

ceratina Ach. (1810)

chaetophora Stirton (1883)

cornuta Körber (1859)

U. inflata (Duby) Motyka (1937)

esperantiana Clerc (1992)

U. fulvoreagens auct. brit. p.p., non (Räsänen) Räsänen (1935)

filipendula Stirton (1881)

flammea Stirton (1881)

florida (L.) Wigg. (1780)

fragilescens Havaas ex Lyngé (1921)

fragilescens var. *mollis* (Vainio) Clerc (1987)
fulvoreagens (Räsänen) Räsänen (1935)
glabrata (Ach.) Vainio (1915)
glabrescens (Nyl. ex Vainio) Vainio (1925)
hirta (L.) Wigg. (1780)
madeirensis Motyka (1964)
rubicunda Stirton (1881)
subfloridana Stirton (1882)
subscabrosa Nyl. ex Motyka (1937)
wasmuthii Räsänen (1932)
wirthii Clerc (1984)

Verrucaria Schrader (1794)

aethiobola Wahlenb. (1803)
V. aquilella Nyl. (1876)
amphibia R. Clem. (1807)
aquatilis Mudd (1861)
aranensis McCarthy (1988)
aspiciliicola R. Sant (1984)
 ? = *V. canella* Nyl. (1883)
V. aspiciliae Zehetl. (1978), non (Lahm) Stizenb. (1882)
 ? *V. glauicina* subsp. *canella* (Nyl.) A.L. Sm. (1911)
baldensis Massal. (1852)
bryoctona (Th. Fr.) A. Orange (1991)
V. psammophila Erichsen (1939)
bulgarica Szat. (1930)
caerulea DC. (1805)
calciseda DC. (1805)
carnea (Arnold) Servít (1948)
cyanea Massal. (1853)
degelii R. Sant. (1939)
ditmarsica Erichsen (1937)
dolosa Hepp (1860)
V. mutabilis auct. p.p., non Borrer ex Leighton (1851)
dufourii DC. (1805)
elaeomeleana (Massal.) Arnold (1868)
funckii (Sprengel) Zahlbr. (1921)
fusconigrescens Nyl. (1872)
V. collematodes auct. brit., non Garov. (1865)
glaucina auct. brit., non Ach. (1810)
halizoa Leighton (1871)
V. mackenzie-lambii Erichsen (1940)
hochstetteri Fr. (1831)
V. dolomitica (Massal.) Krempelh. (1861)
 ? *V. foveolata* (Flörke) Massal. (1852)
hydrela Ach. (1814)
V. hibernica Servít (1955), non Zsch. (1927)
internigrescens (Nyl.) Erichsen (1928)

- knowlesiae* McCarthy (1988)
latebrosa Körber (1855)
latericola Erichsen (1943)
V. granulosaria Clauz. & Zehetl. (1978)
V. insularis auct. brit., non (Massal.) Jatta (1909)
macrostoma Dufour ex DC. (1805)
? *V. griseorubens* Migula (1930)
macrostoma f. *furfuracea* B. de Lesd. (1949)
V. viridula f. *tectorum* auct. brit., non (Massal.) Laundon (1967)
margacea (Wahlenb.) Wahlenb. (1812)
V. leightonii Hepp (1853)
maura Wahlenb. (1803)
mucosa Wahlenb. (1803)
muralis Ach. (1803)
murina Leighton (1851)
nigrescens Pers. (1795)
ochrostoma (Borrer ex Leighton) Trevisan (1860)
papillosa Ach. (1810)
parmigerella Zahlbr. (1919)
pinguicula Massal. (1856)
praetermissa (Trevisan) Anzi (1864)
V. amylacea auct. brit., non Hepp ex Arnold (1858)
prominula Nyl. (1861)
rheitrophila Zsch. (1922)
V. kernstockii Zsch. (1927)
sandstedei B. de Lesd. (1911)
V. lorraine-smithii Knowles (1913)
simplex McCarthy (1988)
striatula Wahlenb. (1803)
viridula (Schrader) Ach. (1803)
xyloxena Norm. (1867)
V. melaenella Vainio (1921)

Vestergrenopsis Gyelnik (1940)
elaeina (Wahlenb. ex Ach.) Gyelnik (1940)

Vezdaea Tscherm.-Woess & Poelt (1976)
acicularis Coppins (1987)
aestivalis (Ohl.) Tscherm.-Woess & Poelt (1976)
leprosa (P. James) Vězda (1977)
retigera Poelt & Döbbeler (1977)
rheocarpa Poelt & Döbbeler (1977)

Vorarlbergia Grumm. (1969) - see *Epigloea*.

Wadeana Coppins & P. James (1978)
dendrographa (Nyl.) Coppins & P. James (1978)
minuta Coppins & P. James (1978)

Xanthoparmelia (Vainio) Hale (1974) - see *Parmelia*.

Xanthoria (Fr.) Th. Fr. (1880)

calcicola Oxner (1937)

X. aureola auct. pro parte, non (Ach.) Erichsen (1928)

candelaria (L.) Th. Fr. (1861)

ectaneoides (Nyl.) Zahlbr. (1931)

X. parietina var. *ectanea* auct., non (Ach.) J.J. Kickx (1867)

elegans (Link) Th. Fr. (1860)

parietina (L.) Th. Fr. (1860)

polycarpa (Hoffm.) Th. Fr. ex Rieber (1891)

ulophyllodes Räsänen (1931)

X. fallax auct. brit., non (Hepp) Arnold (1880)

Xerotrema Sherw. & Coppins (1980) [F]

megalospora Sherw. & Coppins (1980) [F]

Xylographa (Fr.) Fr. (1838)

abietina (Pers.) Zahlbr. (1922)

? *X. laricicola* Nyl. (1875)

trunciseda (Th. Fr.) Minks ex Redinger (1938)

vitiligo (Ach.) Laundon (1963)

Zamenhofia Clauz. & Roux (1985)

coralloidea (P. James) Clauz. & Roux (1985)

Porina coralloidea P. James (1971)

hibernica (P. James & Swinscow) Clauz. & Roux (1985)

Porina hibernica P. James & Swinscow (1962)

rosei (Sérusiaux) P. James (1992)

Porina rosei Sérusiaux (1990)

Acknowledgements

We would like to thank the numerous colleagues both within and outside the British Is. who were involved with the production of the *Flora*, who either discovered new species or helped with nomenclatural aspects.

References

- Cannon, P.F., Hawksworth, D.L. & Sherwood-Pike, M.A. (1985) *The British Ascomycotina: An annotated checklist.* 302 pp. Kew: Commonwealth Mycological Institute.
- Diederich, P. & Sérusiaux, E. (1993) A nomenclatural note on *Lauderlindsaya* (Ascomycotina, Verrucariales). *Lichenologist* **25**: 97-100.
- Egea, J.M. & Torrente, P. (1993) The lichen genus *Bactrospora*. *Lichenologist* **25**: (in ed.).
- Hawksworth, D.L., James, P.W. & Coppins, B.J. (1980) Checklist of British lichen-forming, lichenicolous and allied fungi. *Lichenologist* **12**: 1-115.
- Jørgensen, P.M. (1993) Further notes on the lichen genus *Leptogium* with emphasis on the smaller species. *Lichenologist* **25** (in press).
- Purvis, O.W., Coppins, B.J., Hawksworth, D.L., James, P.W. & Moore, D.M. (1992) *The Lichen Flora of Great Britain and Ireland.* 710 pp. London: The Natural History Museum.
- Stenroos, S. (1989a) Taxonomy of the *Cladonia coccifera* group. 1. *Ann. Bot. Fennic.* **26**: 157-168.
- Stenroos, S. (1989b) Taxonomy of the *Cladonia coccifera* group. 1. *Ann. Bot. Fennic.* **26**: 237-261.
- Thomson, J.W. (1991) The lichen genus *Staurothele* in North America. *Bryologist* **94**: 351-367.
- Torrente, P. & Egea, J.M. (1989) La familia Opegraphaceae en el Área Mediterránea de la Península Ibérica y norte de África. *Bibliotheca lich.* **32**: 1-282.

Appendix 1

Appendix 1 must be treated with caution as it is likely that some may be refound in Britain and it is given to stimulate people to actively search for these species. It is interesting to note that Hawksworth *et al.* (1980) identified 36 species as apparently extinct. Six of these, notably *Arthonia elegans*, *Catapyrenium psoromoides*, *Chaenotheca phaeocephala*, *Lecanora achariana*, *Pertusaria bryontha* and *Usnea glabrata*, have been rediscovered. Furthermore, the material on which a further 8 records are based, notably *Bryoria nitidula*, *Cladonia turgida*, *Phaeophyscia ciliata*, *P. muscigena*, *Ramalina capitata*, *Umbilicaria arctica*, *Usnea cavernosa* and *Varicellaria rhodocarpa*, is now considered to have originated from outside Britain.

Lichens now apparently extinct in Great Britain and Ireland (not seen since 1900)

- Arthonia galactites* (DC.) Dufour (1818)
- A. myriocarpella* Nyl. (1863)
- A. subfuscicola* (Lindsay) Triebel (1991) [LF]
- Arthothelium spectabile* Flotow ex Massal. (1852)
- Aspicilia tuberculosa* (Ach.) Laundon (1986)
- Bacidia polychroa* (Th. Fr.) Körber (1860)
- B. subturgidula* (Nyl.) Zahlbr. (1926)
- Biatora cuprea* (Sommerf.) Fr. (1831)
- Bryoria implexa* (Hoffm.) Brodo & D. Hawksw. (1977)
 - B. pseudofuscescens* (Gyelnik) Brodo & D. Hawksw. (1977)
- Calicium quercinum* Pers. (1797)
- C. trabinellum* (Ach.) Ach. (1803)
- Caloplaca crenulatella* (Nyl.) H. Olivier (1909)
- C. haematites* (Chaub. ex St-Amans) Zwackh (1862)
- C. irribescens* (Arnold) Zahlbr. (1898)
- C. pollinii* (Massal.) Jatta (1900)
- Catillaria minuta* (Massal.) Lettau (1912)
- C. picila* (Massal.) Coppins (1989)
- Cetraria juniperina* (L.) Ach. (1803)
- Chaenothecopsis debilis* (Turner & Borrer ex Sm.) Tibell (1975) [F]
- Cladonia stellaris* (Opiz) Pouzar & Vězda (1971)
- Collema conglomeratum* Hoffm. (1796)
- Cryptothele rhodosticta* (Taylor) Henssen (1990)
- Endocarpon pusillum* Hedwig (1789)
- E. pusillum* var. *pallidum* (Ach.) Körber (1855)
- Lecania coeruleorubella* (Mudd) M. Mayrh. (1988)
- L. fuscella* (Schaerer) Körber (1855)
- L. olivacella* (Nyl.) Zahlbr. (1928)
- Lecanora congesta* Clauz. & Vězda (1969)
- L. fuscescens* (Sommerf.) Nyl. (1871)
 - Lecidea fuscescens* Sommerf. (1824)
- L. populicola* (DC.) Duby (1830)
- L. pruinosa* Chaub. (1821)
- Lecidea advertens* Nyl. (1866)

- L. commaculans Nyl. (1868)
L. paraclitica Nyl. (1872)
Lecidella pulveracea (Schaerer) H. Sydow (1887)
Leptogium hildenbrandii (Garov.) Nyl. (1856)
Melaspilea amota Nyl. (1867)
M. diplasiospora (Nyl.) Müll. Arg. (1887)
M. leciographoides Vouaux (1913) [LF]
Nephroma helveticum Ach. (1810)
N. resupinatum (L.) Ach. (1810)
Pertusaria lactescens Mudd (1861)
P. panyrga auct. brit., non (Ach.) Massal. (1855)
 P. hutchinsiae (Borrer) Leighton (1851)
? Polyblastia gothica Th. Fr. (1865)
Stereocaulon tomentosum Fr. (1824)
Strigula stigmatella (Ach.) R. Harris (1980)
Thelotrema isidioides (Borrer) R. Sant. (1980)
Toninia oribata (Nyl.) P. James (1965)
T. rosulata (Anzi) H. Olivier (1911)
Tornabea scutellifera (With.) Laundon (1984)
 Tornabenia atlantica (Ach.) Kurok. (1962)

Appendix 2

Other excluded species: doubtful taxa or based on dubious records

* = specimen(s) on which record is based collected by J.M. Crombie and probably originating from outside British Isles.

- * *Acarospora peliscypha* Th. Fr. (1860)
- Arthonia patellullata* Nyl. (1853)
- A. punctilliformis* Leighton (1876) [F]
- A. stictarum* (Tul.) Th. Fr. (1874) [LF]
- Arthopyrenia grisea* (Schleicher ex Schaeerer) Körber (1855)
- A. allogena* (Nyl.) A.L. Sm. (1911) [LF]
- A. aphorisa* (Stirton) A.L. Sm. (1911)
- A. desistens* (Nyl.) A.L. Sm. (1911) [LF]
- A. lomnitzensis* B. Stein (1879) [LF]
- A. platypyrenia* (Nyl.) Arnold (1870)
- A. socialis* Körber (1865)
- A. spilobola* (Nyl.) Arnold (1891)
- Aspicilia cupreogrisea* (Th. Fr.) Hue (1912)
- A. verrucigera* Hue (1912)
- Bacidia hemipolia* (Nyl.) Malme (1865)
- B. herbidula* (Nyl.) Zahlbr. (1926)
- B. killiasii* (Hepp) D. Hawksw. (1983) [LF]
- B. microcarpa* (Th. Fr.) Lettau (1912)
- * *Bellemerea cinereorufescens* (Ach.) Clauz. & Roux (1984)
 Aspicilia cinereorufescens (Ach.) Massal. (1852)
- * *Bryocaulon divergens* (Ach.) Kärnefelt (1986)
 Coelocaulon divergens (Ach.) R. Howe (1912)
- * *Bryoria nitidula* (Th. Fr.) Brodo & D. Hawksw. (1977)
- Buellia succedens* (Nyl.) Arnold (1870)
- Caloplaca vitellinula* (Nyl.) H. Olivier (1897)
- * *Candelariella arctica* (Körber) R. Sant. (1966)
- Catillaria erysiboides* (Nyl.) Th. Fr. (1874)
- Cetraria cucullata* (Bellardi) Ach. (1803)
- * *C. odontella* (Ach.) Ach. (1814)
- * *Cladonia acuminata* (Ach.) Norrlin (1875)
- * *C. amăurocraea* (Flörke) Schaeerer (1823)
- C. brevis* (Sandst.) Sandst. (1922)
- * *C. turgida* Hoffm. (1796)
- Collema ryssoleum* (Tuck.) A. Schneider (1898)
- Cryptothele granuliformis* (Nyl.) Henssen (1984)
 Pyrenopsisidium granuliforme (Nyl.) Forss. (1885)
- Epigloea renitens* (Grumm.) Döbbeler (1984)
- * *Hypogymnia vittata* (Ach.) Parrique (1898)
- Lecania dubitans* (Nyl.) A. L. Sm. (1928)
- Lecanora cenisia* var. *atrynea* (Ach.) Clauz. & Roux (1985)
- L. pacifica* Tuck. (1882)
- Lecidea callista* Stirton (1874)

- L. calpodes* Stirton (1876)
L. sincerula Nyl. (1876)
L. stirtoniana Zahlbr. (1925)
L. relicta Stirton (1876)
Lecidella euphorea (Flörke) Hertel (1990)
Lecidea dolosa (Nyl.) Vainio (1893)
Lempholemma isidioides (Nyl. ex Arnold) Magnusson (1939)
Leptogium minutissimum (Flörke) Fr. (1846)
Lithographa tesserata var. *petraea* auct. brit., non (Ach.) Redinger (1937)
Melaspilea constrictella (Stirton) A.L. Sm. (1911)
Micarea melaenida (Nyl.) Coppins (1983)
Catillaria melaenida (Nyl.) Arnold (1870)
Mycomicrothelia mélanospora (Hepp) D. Hawksw. (1982)
Mycóporum quercus var. *ilicis* (Riedl) Coppins (1979) [F]
* *Ochrolechia upsaliensis* (L.) Massal. (1852)
* *Ophioparma lapponica* (Räsänen) Hafellner & R.W. Rogers (1988)
Haematomma lapponica Räsänen (1931)
Parmelia afrorevoluta Krog & Swinscow (1979)
P. centrifuga (L.) Ach. (1803)
Arctoparmelia centrifuga (L.) Hale (1986)
* *P. olivacea* (L.) Ach. (1803)
Peltigera aphthosa (L.) Willd. (1787)
Peltula euploca (Ach.) Poelt ex Ozenda & Clauz. (1970)
* *Perthusaria carneopallida* (Nyl.) Anzi ex Nyl. (1868)
P. trochiscea Norman (1868)
* *Phaeophyscia ciliata* (Hoffm.) Moberg (1977)
* *Physcia phaea* (Tuck.) Thompson (1963)
* *Physconia muscigena* (Ach.) Poelt (1965)
Placynthium majus Harm. (1913)
* *Ramalina capitata* (Ach.) Nyl. (1879)
* *R. dilacerata* (Hoffm.) Hoffm (1825)
* *R. thrausta* (Ach.) Nyl. (1860)
Rhizocarpon grande (Flörke ex Flotow.) Arnold (1871)
R. roridulum (Th. Fr.) H. Olivier (1981)
R. transiens Eitner (1911)
Rinodina arnoldii Mayrh. & Poelt (1979)
R. trachyticá (Massal.) Bagl. & Car. (1880)
Stereocaulon botryosum Ach. (1810)
S. paschale (L.) Hoffm. (1796)
Strigula affinis (Massal.) R. Harris (1980)
Thelidium cataractarum (Hepp) Lönnr. (1859)
T. minimum (Massal. ex Körber) Arnold (1871)
* *Tonina alutacea* (Anzi) H. Olivier (1911)
* *T. candida* (Web. ex Wigg.) Th. Fr. (1867)
T. tristis (Th. Fr.) Th. Fr. (1874)
* *Umbilicaria arctica* (Ach.) Nyl. (1859)
* *U. leiocarpa* DC. in Lam. & DC. (1805)
* *Usnea cavernosa* (Tuck.) (1850)

- U. foveata Vainio (1928)
- U. lapponica Vainio (1925)
- * Varicellaria rhodocarpa (Körber) Th. Fr. (1871)
- Verrucaria conturmatula Nyl. (1879) [LF]
- V. phaeosperma Arnold (1874)
- V. pseudomemnonia Zsch. (1924)

Appendix 3

Species newly recorded from Great Britain and Ireland

Included are records of lichens recognized as occurring in Great Britain and Ireland subsequent to publication of the *Flora*.

- Acarospora durietzii Magnusson (1924)
- Arthonia abelonae P.M. Jørg. (1970) [LF]
- Bacidia igniari (Nyl.) Oxner (1968)
- B. myrtillilcola Erichsen (1939)
- Carbonea assimilis (Körber) Hafellner & Hertel (1987)
- Geltingia groenlandiae Alstrup & D. Hawksw. (1990) [LF]
- Lecanora mughicola Nyl. (1872)
- L. persimilis (Th. Fr.) Nyl. (1876)
- Lepraria jackii Tønsb. (1992)
- * Leptogium byssinum (Hoffm.) Zwackh ex Nyl. (1856)
- L. imbricatum P.M. Jørg. in ed. (1993)
- L. intermedium (Arnold) Arnold (1885)
- Lichenodium sirosiphoides Nyl. (1856)
- Lopadium coralloideum (Nyl.) Lynge (1940)
- Opegrapha ? monspeliensis Nyl. (1857)
- O. xerica Torrente & Egea (1992)
- Parmelia protomatrae Gyelnik (1931)
- Pyrenula ? microtheca R. Harris (1989)
- Rhizocarpon ochrolechia (Poelt & Nimis) Hafellner (1992)
- Rimularia sphacelata (Th. Fr.) Hertel & Rambold (1990)
- Sarcopyrenia cylindrospora (P. Crouan & H. Crouan) M.B. Aguirre (1990)
- Thelidium aeneovinosum Anzi (Arnold) (1868)
- Verrucaria carnea (Arnold) Servit (1948)

* material previously referred to this species in Britain = *L. biatorinum*
(Nyl.) Leighton

Appendix 4

New combinations.

Biatora gyrophorica (Tønsb.) Coppins comb. nov. Bas.: *Lecidea gyrophorica* Tønsb., *Sommerfeltia* 14: 172 (1992).

Cladonia polydactyla var. **umbricola** (Tønsb. & Ahti) Coppins comb. nov.
Bas.: *Cladonia umbricola* Tønsb. & Ahti, *Norw. J. Bot.* 27: 307 (1980).

Moelleropsis humida (Kullh.) Coppins & P.M. Jørg. comb. nov. Bas.:
Biatora humida Kullh., *Not. Salsk. Fauna Fl. fenn. Förh.* 11: 274 (1871).

Rimularia intercedens (Magnusson) Coppins comb. nov. Bas.: *Lecanora intercedens* Magnusson, *Acta Horti Gothob.* 6: 17-19 (1930) ("1931").

Rimularia mullensis (Stirton) Coppins comb. nov. Bas.: *Lecidea mullensis* Stirton, *Scott. Nat.* 4: 166 (1877).

Notes

Notes

Notes

BULLETIN 72 (Supplement). Issued by the British Lichen Society
(Registered Charity No 228850), c/o Department of Botany, Natural
History Museum , Cromwell Road, London, SW7 5BD (Telephone 071938
8852).

Printed by DESA Ltd (Nottingham).

ISSN 0300 - 4562

